

On automorphic sheaves on Bun_G

Sergey Lysenko

ABSTRACT Let X be a smooth projective connected curve over an algebraically closed field k of positive characteristic. Let G be a reductive group over k , γ be a dominant coweight for G , and E be an ℓ -adic $\check{G}$ -local system on X , where $\check{G}$ denotes the Langlands dual group (over $\bar{\mathbb{Q}}_\ell$). Let Bun_G be the moduli stack of G -bundles on X .

Under some conditions on the triple (G, γ, E) we propose a conjectural construction of a distinguished E -Hecke automorphic sheaf on Bun_G . We are motivated by a construction of automorphic forms suggested by Ginzburg, Rallis and Soudry in [6, 7]. We also generalize Laumon's theorem ([10], Theorem 4.1) for our setting. Finally, we formulate an analog of the Vanishing Conjecture of Frenkel, Gaitsgory and Vilonen for Levi subgroups of G .

1. INTRODUCTION

Let F be a number field, $\mathbb{A}$ be its ring of adeles. If G is one of the groups SO_{2n+1} , Sp_{2n} or SO_{2n} then consider standard representation $\check{G} \rightarrow \check{H}$ of the Langlands dual group $\check{G}$, so here H is GL_{2n} , GL_{2n+1} or GL_{2n} respectively. For an irreducible, automorphic, cuspidal representation τ of $H(\mathbb{A})$ satisfying some additional conditions, D. Ginzburg, S. Rallis and D. Soudry have proposed a conjectural construction of an irreducible, automorphic cuspidal representation σ of $G(\mathbb{A})$ which lifts to τ (cf., [6, 7]).

For example, consider $G = \mathrm{SO}_{2n+1}$. Let X be a smooth projective absolutely irreducible curve over $\mathbb{F}_q$. Consider the Langlands dual group $\check{G} = \mathrm{Sp}_{2n}$ over $\bar{\mathbb{Q}}_\ell$. Let $H = \mathrm{GL}_{2n}$ over $\mathbb{F}_q$, $\check{H} = \mathrm{GL}_{2n}$ over $\bar{\mathbb{Q}}_\ell$. The standard representation V of $\check{G}$ is a map $\check{G} \rightarrow \check{H} = \mathrm{GL}(V)$.

Let E be an ℓ -adic $\check{G}$ -local system on X , assume that V_E is irreducible. According to [8], theorem VII.6, irreducibility of V_E implies that $\mathrm{End}(V_E)$ is pure of weight zero. It follows that for each closed point $x \in X$ the local L-function $L(E_x, \check{\mathfrak{g}}, s)$ is regular at $s = 1$, and the corresponding irreducible unramified representation of $G(F_x)$ is generic. Here F_x denotes the completion of $\mathbb{F}_q(X)$ at x , and $\check{\mathfrak{g}} = \mathrm{Lie} \check{G}$. An analog of D. Ginzburg, S. Rallis and D. Soudry's conjecture for function field is to predict that in the L-packet of automorphic forms corresponding to E there exists a unique nonramified cuspidal generic form $\varphi_E : \mathrm{Bun}_G(\mathbb{F}_q) \rightarrow \bar{\mathbb{Q}}_\ell$ (cf. [7], Conjecture on p. 809 and [6]).

In *loc.cit.* an additional condition is required: the L-function $L(E, \wedge^2 V, s)$ has a pole of order exactly one at $s = 1$. This condition is satisfied in our situation. Indeed, $\wedge^2 V = V' \oplus \bar{\mathbb{Q}}_\ell$, where V' is an irreducible representation of $\check{G}$. Since V is self-dual, $H^0(X \otimes \bar{\mathbb{F}}_q, V'_E) = 0$ and the L-function $L(E, V', s)$ is a polynomial in q^{-s} . The purity argument shows that $L(E, V', 1) \neq 0$.

In this paper we consider the problem of constructing a geometric counterpart of φ_E . Given a reductive group G , a dominant coweight γ and a $\check{G}$ -local system E on X , we impose on these data some conditions similar to the above. Then we propose a conjectural construction of a

distinguished E -Hecke eigensheaf on the moduli stack Bun_G of G -bundles on X . Our approach applies to root systems A_n, B_n, C_n for all n , D_n for odd n , and also E_6, E_7 . For GL_n our method reduces to the one proposed by Laumon in [9].

The construction is exposed in Sect. 2, 3. In Sect. 4 we study the additional structure on Levi subgroups induced by γ , and prove a generalization of Laumon's theorem ([10], Theorem 4.1) for our setting. We discuss its applications to cuspidality and formulate an analog of the Vanishing Conjecture of Frenkel, Gaitsgory and Vilonen for Levi subgroups of G .

2. STATEMENTS AND CONJECTURES

2.1 NOTATION Throughout, k will denote an algebraically closed field of characteristic $p > 0$. Let X be a smooth projective connected curve over k . Fix a prime $\ell \neq p$. For a k -scheme (or k -stack) S write $\text{D}(S)$ for the bounded derived category of ℓ -adic étale sheaves on S .

Let G be a connected reductive group over k . Fix a Borel subgroup $B \subset G$. Let $N \subset B$ be its unipotent radical and $T = B/N$ be the "abstract" Cartan. Let Λ denote the coweight lattice. The weight lattice is denoted by $\check{\Lambda}$. The semigroup of dominant coweights (resp., weights) is denoted Λ^+ (resp., $\check{\Lambda}^+$). The set of vertices of the Dynkin diagram of G is denoted by $\mathcal{I}$. To each $i \in \mathcal{I}$ there corresponds a simple root $\check{\alpha}_i$ and a simple coroot α_i . By $\check{\rho} \in \check{\Lambda}$ is denoted the half sum of positive roots of G and by w_0 the longest element of the Weil group W .

Let Λ^{pos} denote $\mathbb{Z}_+$ -span of positive coroots. The set Λ^+ is equipped with the order $\nu_1 \leq \nu_2$ iff $\nu_2 - \nu_1 \in \Lambda^{pos}$. Similarly, we have an order on $\check{\Lambda}^+$.

To a dominant weight $\check{\lambda}$ one attaches the Weil G -module $\mathcal{V}^{\check{\lambda}}$ with a fixed highest weight vector $v^{\check{\lambda}} \in \mathcal{V}^{\check{\lambda}}$. For any pair $\check{\lambda}, \check{\nu} \in \check{\Lambda}^+$ there is a canonical map $\mathcal{V}^{\check{\lambda}+\check{\nu}} \rightarrow \mathcal{V}^{\check{\lambda}} \otimes \mathcal{V}^{\check{\nu}}$ sending $v^{\check{\lambda}+\check{\nu}}$ to $v^{\check{\lambda}} \otimes v^{\check{\nu}}$.

For $\lambda \in \Lambda^+$ write V^λ for the irreducible representation of $\check{G}$ of highest weight λ .

The trivial G -bundle on a scheme is denoted by $\mathcal{F}_G^0$. Recall that for any finite subfield $k' \subset k$ and any non-trivial character $\psi : k' \rightarrow \mathbb{Q}_\ell$ one can construct the Artin-Shrier sheaf $\mathcal{L}_\psi$ on $\mathbb{G}_{a,k}$. The intersection cohomology sheaves are normalized to be pure of weight zero.

2.2 ADDITIONAL DATA AND ASSUMPTIONS We say that $\gamma \in \Lambda^+$ is *minuscule* if γ is a minimal element of Λ^+ and $\gamma \neq 0$. If $\gamma \in \Lambda^+$ is minuscule then, by (Lemma 1.1, [14]), for any root $\check{\alpha}$ we have $\langle \gamma, \check{\alpha} \rangle \in \{0, \pm 1\}$, and the set of weights of V^γ coincides with the W -orbit of γ . For example, if $\gamma \neq 0$ is orthogonal to all roots then γ is minuscule. One checks that the natural map from the set of minuscule dominant coweights to $\pi_1(G)$ is injective.

Definition 1. We say that $\{\gamma\}$ is a *1-admissible datum* if the following conditions hold

- the center $Z(G)$ is a connected 1-dimensional torus;
- $\pi_1(G) \xrightarrow{\sim} \mathbb{Z}$;
- $\gamma \in \Lambda^+$ is a minuscule dominant coweight whose image θ in $\pi_1(G)$ generates $\pi_1(G)$;
- V^γ is a faithful representation of $\check{G}$.

Fix a 1-admissible datum γ . For $k \geq 0$ set $\Lambda_{G,S}^{+,k\theta} = \{\mu \in \Lambda^+ \mid \mu \leq k\gamma\}$. Let $\Lambda_{G,S}^+$ be the union of $\Lambda_{G,S}^{+,k\theta}$ for $k \geq 0$. Set

$$\check{\Lambda}_S^+ = \{\check{\lambda} \in \check{\Lambda}^+ \mid \langle w_0(\lambda), \check{\lambda} \rangle \geq 0 \text{ for any } \lambda \in \Lambda_{G,S}^+\}$$

Let $\check{\omega}_0$ be the generator of the group of weights orthogonal to all coroots, we fix $\check{\omega}_0$ by requiring $\langle \theta, \check{\omega}_0 \rangle = 1$. For $i \in \mathcal{I}$ denote by $\check{\omega}_i \in \check{\Lambda}^+$ the fundamental weight corresponding to α_i that satisfies $\langle w_0(\gamma), \check{\omega}_i \rangle = 0$. Note that $\check{\omega}_0, \check{\omega}_i (i \in \mathcal{I})$ form a basis of $\check{\Lambda}$.

The following lemma is straightforward.

Lemma 1. *The semigroup $\check{\Lambda}_S^+$ is the $\mathbb{Z}_+$ -span of $\check{\omega}_0, \check{\omega}_i (i \in \mathcal{I})$. Besides*

$$\Lambda_{G,S}^+ = \{\lambda \in \Lambda^+ \mid \langle w_0(\lambda), \check{\lambda} \rangle \geq 0 \text{ for all } \check{\lambda} \in \check{\Lambda}_S^+\} \quad \square$$

Since $\pi_1(G) \xrightarrow{\sim} \mathbb{Z}$, it follows that $[G, G]$ is simply-connected. Note that for $\check{\mu} \in \check{\Lambda}^+, \check{\lambda} \in \check{\Lambda}_S^+$ the condition $\check{\mu} \leq \check{\lambda}$ implies $\check{\mu} \in \check{\Lambda}_S^+$. Note that $\{-w_0(\gamma)\}$ is also a 1-admissible datum.

Since V^γ is faithful, the weights of V^γ generate Λ and for each $i \in \mathcal{I}$ we have $\langle \gamma, \check{\omega}_i \rangle > 0$. For each maximal positive root $\check{\alpha}$ we have $\langle \gamma, \check{\alpha} \rangle = 1$. In particular, if the root system of G is irreducible (so, nonempty) then γ is a fundamental coweight corresponding to some simple root.

Some examples of 1-admissible data are given in the appendix.

2.2.1 Consider the formal disk $\mathcal{D} = \mathrm{Spf}(k[[t]])$. Recall that the Affine Grassmanian Gr_G is the ind-scheme classifying pairs $(\mathcal{F}_G, \beta)$, where $\mathcal{F}_G$ is a G -bundle on $\mathcal{D}$ and $\beta : \mathcal{F}_G \xrightarrow{\sim} \mathcal{F}_G^0$ is a trivialization over the punctured disk $\mathcal{D}^* = \mathrm{Spec} k((t))$. Define the positive part $\mathrm{Gr}_G^+ \subset \mathrm{Gr}_G$ of Gr_G as a closed subscheme given by the following condition:

$\mathcal{F}_G \in \mathrm{Gr}_G^+$ if for every $\check{\lambda} \in \check{\Lambda}_S^+$ the map

$$\beta^{\check{\lambda}} : \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}|_{\mathcal{D}^*} \rightarrow \mathcal{V}_{\mathcal{F}_G^0}^{\check{\lambda}}|_{\mathcal{D}^*}$$

is regular on $\mathcal{D}$. Note that Gr_G^+ is invariant under the natural action of $G(k[[t]])$.

Recall that for $\mu \in \Lambda^+$ one has the closed subscheme $\overline{\mathrm{Gr}}_G^\mu \subset \mathrm{Gr}_G$ (cf. [2], sect. 3.2). One checks that $\overline{\mathrm{Gr}}_G^\mu \subset \mathrm{Gr}_G^+$ iff $\mu \in \Lambda_{G,S}^+$. Let $\pi_1^+(G) \subset \pi_1(G)$ be the image of $\Lambda_{G,S}^+$ under the projection $\Lambda \rightarrow \pi_1(G)$.

For $\nu \in \pi_1(G)$ the connected component Gr_G^ν of Gr_G is given by the condition:

$$\mathcal{V}_{\mathcal{F}_G^0}^{\check{\omega}_0}(-\langle \nu, \check{\omega}_0 \rangle) \xrightarrow{\sim} \mathcal{V}_{\mathcal{F}_G}^{\check{\omega}_0}$$

For $\nu \in \pi_1(G)$ set $\mathrm{Gr}_G^{+, \nu} = \mathrm{Gr}_G^+ \cap \mathrm{Gr}_G^\nu$.

2.3 Denote by Bun_G the moduli stack of G -bundles on X . Let $\mathcal{H}_G^+$ be the corresponding positive part of the Hecke stack, it classifies collections: $\mathcal{F}_G, \mathcal{F}'_G \in \mathrm{Bun}_G$, an effective divisor D on X , an isomorphism $\beta : \mathcal{F}_G|_{X-D} \xrightarrow{\sim} \mathcal{F}'_G|_{X-D}$ such that for each $\check{\lambda} \in \check{\Lambda}_S^+$ the map

$$\beta^{\check{\lambda}} : \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}'_G}^{\check{\lambda}}$$

extends to an inclusion of coherent sheaves on X , and $\mathcal{V}_{\mathcal{F}_G}^{\check{\omega}_0}(D) \xrightarrow{\sim} \mathcal{V}_{\mathcal{F}'_G}^{\check{\omega}_0}$. For $k \geq 0$ let $\mathcal{H}_G^{+,k} \subset \mathcal{H}_G^+$ be given by $\deg D = k$.

2.4 VERSION OF LAUMON'S SHEAF Given a local system W on X and $d \geq 0$, one defines a sheaf $\mathcal{L}_W^d$ on $\mathcal{H}_G^{+,d}$ as follows.

Let $\tilde{\mathcal{H}}_G^{+,d}$ be the stack of collections: $(\mathcal{F}_G^1, \dots, \mathcal{F}_G^{d+1})$ and $(\beta^i, x_i \in X)_{i=1, \dots, d}$, where $\mathcal{F}_G^i \in \text{Bun}_G$ and

$$\beta^i : \mathcal{F}_G^i|_{X-x_i} \xrightarrow{\sim} \mathcal{F}_G^{i+1}|_{X-x_i}$$

is an isomorphism such that $(\mathcal{F}_G^i, \mathcal{F}_G^{i+1}, \beta^i, x_i) \in \mathcal{H}_G^{+,1}$ for $i = 1, \dots, d$.

Note that $\tilde{\mathcal{H}}_G^{+,d}$ is smooth, because γ is minuscule. We have a convolution diagram

$$\begin{array}{ccc} \tilde{\mathcal{H}}_G^{+,d} & \xrightarrow{\widetilde{\text{supp}}} & X^d \\ \downarrow p & & \downarrow \\ \mathcal{H}_G^{+,d} & \xrightarrow{\text{supp}} & X^{(d)} \end{array}$$

where $\widetilde{\text{supp}}$ (resp., p) sends the above collection to $(x_1, \dots, x_d)$ (resp., to $(\mathcal{F}_G^1, \mathcal{F}_G^{d+1}, \beta, D)$ with $D = x_1 + \dots + x_d$). Let ${}^{rss}X^{(d)} \subset X^{(d)}$ be the open subscheme classifying reduced divisors. Over ${}^{rss}X^{(d)}$, this diagram is cartesian.

The following proposition is an immediate corollary of (Lemma 9.3 [14]).

Proposition 1. *The map p is representable proper surjective and small. $\square$*

Set $\mathcal{S}pr_W^d = p(\widetilde{\text{supp}}^* W^{\boxtimes d}[m](\frac{m}{2}))$ with $m = \dim \mathcal{H}_G^{+,d}$. This is a perverse sheaf, the Goresky-MacPherson extension from $\text{supp}^{-1}({}^{rss}X^{(d)})$. It is equipped with a canonical action of S_d . Define $\mathcal{L}_W^d$ to be the S_d -invariants of $\mathcal{S}pr_W^d$. We have $\mathbb{D}(\mathcal{S}pr_W^d) \xrightarrow{\sim} \mathcal{S}pr_{W*}^d$ canonically and $\mathbb{D}(\mathcal{L}_W^d) \xrightarrow{\sim} \mathcal{L}_{W*}^d$.

We have a diagram

$$\text{Bun}_G \xleftarrow{\mathfrak{p}} \mathcal{H}_G^+ \xrightarrow{\mathfrak{q}} \text{Bun}_G$$

where $\mathfrak{p}$ (resp., $\mathfrak{q}$) sends $(\mathcal{F}_G, \mathcal{F}'_G, \beta)$ to $\mathcal{F}_G$ (resp., $\mathcal{F}'_G$). By (property 3, sect. 5.1.2 [2]), the sheaf $\mathcal{L}_W^d$ is ULA with respect to both projections $\mathfrak{p}$ and $\mathfrak{q}$.

Let $r = \dim V^\gamma$. For a partition $\mu = (\mu_1 \geq \dots \geq \mu_r \geq 0)$ of d define the polynomial functor $W \mapsto W_\mu$ of a $\bar{\mathbb{Q}}_\ell$ -vector space W by

$$W_\mu = (W^{\otimes d} \otimes U_\mu)^{S_d},$$

where U_μ stands for the irreducible representation of S_d corresponding to μ . For $d > 0$ let $\ell(\mu)$ be the greatest index $i \leq r$ such that $\mu_i \neq 0$. For $d = 0$ let $\ell(\mu) = 0$. If $\ell(\mu)$ is less or equal to $\dim W$ then W_μ is the irreducible representation of $\text{GL}(W)$ with h.w. μ , otherwise it vanishes.

For $\nu \in \Lambda^+$ let $\mathcal{A}_\nu$ denote the IC-sheaf on $\overline{\text{Gr}}^\nu$. Recall that the category $\text{Sph}(\text{Gr}_G)$ of spherical perverse sheaves on Gr_G consists of direct sums of $\mathcal{A}_\nu$, as ν ranges over the set of dominant coweights. We have the Satake equivalence of tensor categories $\text{Loc} : \text{Rep}(\check{G}) \rightarrow \text{Sph}(\text{Gr}_G)$ (cf. Theorem 3.2.8, [2]). In particular, we have $\text{Loc}(V^\nu) = \mathcal{A}_\nu$.

Consider a k -point $(\sum d_k x_k, \mathcal{F}'_G)$ of $X^{(d)} \times \text{Bun}_G$. The fibre of $\text{supp} \times \mathfrak{q} : \mathcal{H}_G^{+,d} \rightarrow X^{(d)} \times \text{Bun}_G$ over this point identifies with

$$\prod_k \overline{\text{Gr}}_G^{d_k \gamma} \quad (1)$$

Set $d_G = \dim \text{Bun}_G$. The Satake equivalence yields the following description.

Proposition 2. *For any local system W on X the restriction of $\mathcal{L}_W^d$ to the fibre (1) of $\text{supp} \times \mathfrak{q}$ identifies with the exterior product $(\boxtimes_k \mathcal{L}_k)[d + d_G](\frac{d+d_G}{2})$, where each $\mathcal{L}_k$ is*

$$\mathcal{L}_k = \bigoplus_{\mu} \text{Loc}((V^{\gamma})_{\mu}) \otimes (W_{x_k})_{\mu},$$

the sum being taken over the set of partitions of d_k of length $\leq r$. $\square$

2.5 Given a local system W on X , for $d \geq 0$ define a functor $\text{Av}_W^d : \text{D}(\text{Bun}_G) \rightarrow \text{D}(\text{Bun}_G)$ by

$$\text{Av}_W^d(K) = \mathfrak{q}_!(\mathfrak{p}^* K \otimes \mathcal{L}_W^d)[-d_G](\frac{-d_G}{2})$$

Let also $\text{Av}_W^{-d} : \text{D}(\text{Bun}_G) \rightarrow \text{D}(\text{Bun}_G)$ be given by

$$\text{Av}_W^{-d}(K) = \mathfrak{p}_!(\mathfrak{q}^* K \otimes \mathcal{L}_W^d)[-d_G](\frac{-d_G}{2})$$

The functors Av_W^d and Av_W^{-d} are both left and right adjoint to each other. As in (Proposition 9.5, [4]) one proves

Proposition 3. *Let K be a Hecke eigensheaf on Bun_G with respect to a $\check{G}$ -local system E . Then for the diagram*

$$\text{Bun}_G \times X^{(d)} \xleftarrow{\mathfrak{p} \times \text{supp}} \mathcal{H}_G^{+,d} \xrightarrow{\mathfrak{q}} \text{Bun}_G$$

and any local system W on X we have

$$(\mathfrak{p} \times \text{supp})_!(\mathfrak{q}^* K \otimes \mathcal{L}_W^d)[-d_G](\frac{-d_G}{2}) \xrightarrow{\sim} K \boxtimes (W \otimes V_E^{\gamma})^{(d)}[d](\frac{d}{2}) \quad \square$$

2.6.1 For a T -torsor $\mathcal{F}_T$ on X denote by $\overline{\text{Bun}}_N^{\mathcal{F}_T}$ the stack of collections $(\mathcal{F}_G, \kappa)$, where $\mathcal{F}_G \in \text{Bun}_G$ and for $\check{\lambda} \in \check{\Lambda}^+$

$$\kappa^{\check{\lambda}} : \mathcal{L}_{\mathcal{F}_T}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}$$

are inclusions of coherent sheaves on X satisfying Plücker relations (as in [3], section 2.1.2).

The open substack $j : \text{Bun}_N^{\mathcal{F}_T} \hookrightarrow \overline{\text{Bun}}_N^{\mathcal{F}_T}$ is given by the condition that all $\kappa^{\check{\lambda}}$ are maximal embeddings.

Consider the stack of pairs $(\mathcal{F}_T, \tilde{\omega})$, where $\mathcal{F}_T$ is a T -torsor on X and $\tilde{\omega}$ is a trivial conductor, that is, $\tilde{\omega}$ is a collection of isomorphisms $\tilde{\omega}_i : \mathcal{L}_{\mathcal{F}_T}^{\check{\alpha}_i} \xrightarrow{\sim} \Omega$ for each $i \in \mathcal{I}$. The exact sequence $1 \rightarrow Z(G) \rightarrow T \rightarrow \prod_{i \in \mathcal{I}} \mathbb{G}_m \rightarrow 1$, where the second map is $\prod_{i \in \mathcal{I}} \check{\alpha}_i$, shows that this stack is noncanonically isomorphic to $\text{Bun}_{Z(G)}$ (recall that by our assumption $Z(G)$ is connected).

Fix a section $T \rightarrow B$. Then for each pair $(\mathcal{F}_T, \tilde{\omega})$ we have the evaluation map $\text{ev}^{\tilde{\omega}} : \text{Bun}_N^{\mathcal{F}_T} \rightarrow \mathbb{A}^1$ (cf. [3], section 4.1.1). Fix a T -torsor on X with a trivial conductor $(\mathcal{F}_T, \tilde{\omega})$.

Remark 1. If $(\mathcal{F}'_T, \tilde{\omega}')$ is another T -torsor with trivial conductor on X then there exists a $Z(G)$ -torsor $\mathcal{F}_{Z(G)}$ on X and an isomorphism $\mathcal{F}_T \otimes \mathcal{F}_{Z(G)} \xrightarrow{\sim} \mathcal{F}'_T$ with the following property. Let $\text{Bun}_N^{\mathcal{F}_T} \rightarrow \text{Bun}_N^{\mathcal{F}'_T}$ be the isomorphism that sends $\mathcal{F}_B$ to $(\mathcal{F}_B \times \mathcal{F}_{Z(G)})/Z(G)$, where $Z(G)$ acts diagonally. Then the diagram commutes

$$\begin{array}{ccc} \text{Bun}_N^{\mathcal{F}_T} & \xrightarrow{\sim} & \text{Bun}_N^{\mathcal{F}'_T} \\ & \searrow \text{ev}^{\tilde{\omega}} & \downarrow \text{ev}^{\tilde{\omega}'} \\ & & \mathbb{A}^1 \end{array}$$

2.6.2 For $d \geq 0$ let $\mathcal{Y}_d$ be the stack of collections $(\mathcal{F}_G, D \in X^{(d)}, \kappa)$, where $(\mathcal{F}_G, \kappa) \in \overline{\text{Bun}}_N^{\mathcal{F}'_T}$ with $\mathcal{F}'_T = \mathcal{F}_T(w_0(\gamma)D)$. So, for each $\check{\lambda} \in \check{\Lambda}_S^+$ we have an embedding of coherent sheaves

$$\kappa^{\check{\lambda}} : \mathcal{L}_{\mathcal{F}_T}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}},$$

and $\kappa^{\tilde{\omega}_0}$ induces an isomorphism $\mathcal{L}_{\mathcal{F}_T}^{\tilde{\omega}_0}(D) \xrightarrow{\sim} \mathcal{V}_{\mathcal{F}_G}^{\tilde{\omega}_0}$.

Let ${}_0\mathcal{Y}_d \subset \mathcal{Y}_d$ denote the open substack given by the condition $(\mathcal{F}_G, \kappa) \in \text{Bun}_N^{\mathcal{F}'_T}$.

Consider the fibred product $\mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,k}$, where the map $\mathcal{H}_G^{+,k} \rightarrow \text{Bun}_G$ is $\mathfrak{p}$. For $k \geq 0$ we have a proper representable map

$$\mathfrak{q}_Y : \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,k} \rightarrow \mathcal{Y}_{d+k}$$

that sends $(\mathcal{F}_G, \kappa, \mathcal{F}'_G, \beta : \mathcal{F}_G|_{X-D'} \xrightarrow{\sim} \mathcal{F}'_G|_{X-D'})$ to $(\mathcal{F}'_G, \kappa')$, where $\kappa'^{\check{\lambda}}$ are the compositions

$$\mathcal{L}_{\mathcal{F}_T}^{\check{\lambda}} \xrightarrow{\kappa^{\check{\lambda}}} \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}'_G}^{\check{\lambda}}$$

Given a local system W on X , define the sheaf $\mathcal{P}_{W,\psi}^d$ on $\mathcal{Y}_d$ as follows. Consider the open immersion $j : \text{Bun}_N^{\mathcal{F}_T} \rightarrow \overline{\text{Bun}}_N^{\mathcal{F}_T} \xrightarrow{\sim} \mathcal{Y}_0$. Set

$$\mathcal{P}_{W,\psi}^0 = \mathcal{P}_\psi^0 = j_!(\text{ev}^{\tilde{\omega}})^* \mathcal{L}_\psi[d_N]\left(\frac{d_N}{2}\right),$$

where $d_N = \dim \text{Bun}_N^{\mathcal{F}_T}$. By (Theorem 2, [3]), this is a perverse sheaf and $\mathbb{D}(\mathcal{P}_\psi^0) \xrightarrow{\sim} \mathcal{P}_{\psi^{-1}}^0$. For $d > 0$ set

$$\mathcal{P}_{W,\psi}^d = \mathfrak{q}_{Y!}(\mathcal{P}_\psi^0 \boxtimes \mathcal{L}_W^d)[-d_G]\left(\frac{-d_G}{2}\right)$$

It is easy to see that $\bar{\mathbb{Q}}_\ell \boxtimes \mathcal{L}_W^d$ is ULA with respect to the projection $\mathcal{Y}_0 \times_{\text{Bun}_G} \mathcal{H}_G^{d,+} \rightarrow \mathcal{Y}_0$. So, by (property 5, sect. 5.1.2 [2]),

$$\mathbb{D}(\mathcal{P}_\psi^0 \boxtimes \mathcal{L}_W^d)[-d_G]\left(\frac{-d_G}{2}\right) \xrightarrow{\sim} \mathcal{P}_{\psi^{-1}}^0 \boxtimes \mathcal{L}_{W^*}^d[-d_G]\left(\frac{-d_G}{2}\right)$$

Therefore, $\mathbb{D}(\mathcal{P}_{W,\psi}^d) \xrightarrow{\sim} \mathcal{P}_{W^*,\psi^{-1}}^d$ canonically.

Let ${}^{rss}\mathcal{Y}_d \subset {}_0\mathcal{Y}_d$ be the preimage of ${}^{rss}X^{(d)}$ under the projection ${}_0\mathcal{Y}_d \rightarrow X^{(d)}$.

Proposition 4. *For any local system W on X , $\mathcal{P}_{W,\psi}^d$ is a perverse sheaf on $\mathcal{Y}_d$, the Goresky-MacPherson extension from ${}^{rss}\mathcal{Y}_d$. If W is irreducible then $\mathcal{P}_{W,\psi}^d$ is irreducible.*

The proof is found in Section 3.2.

2.7 Let $\pi^0 : \text{Bun}_N^{\mathcal{F}_T} \rightarrow \text{Bun}_G$ be the projection.

Definition 2. Let K be a E -Hecke eigensheaf on Bun_G , where E is a $\check{G}$ -local system on X . We say that K is *generic normalized* if it is equipped with an isomorphism

$$\text{R}\Gamma_c(\text{Bun}_N^{\mathcal{F}_T}, \mathcal{P}_\psi^0 \otimes \pi^{0*}K) \xrightarrow{\sim} \bar{\mathbb{Q}}_\ell[d_G](\frac{d_G}{2})$$

By Remark 1, this property does not depend (up to a tensoring K by a 1-dimensional vector space) on our choice of the pair $(\mathcal{F}_T, \tilde{\omega})$.

Write Bun_G^d for the connected component of Bun_G given by $\deg \mathcal{F}_G = \deg \mathcal{F}_T + d\theta$. Let $\pi : \mathcal{Y}_d \rightarrow \text{Bun}_G^d$ and $\phi : \mathcal{Y}_d \rightarrow X^{(d)}$ be the projections. From Proposition 3 one derives

Corollary 1. *Let K be a generic normalized E -Hecke eigensheaf on Bun_G . Let W be any local system on X . Then for each $d \geq 0$ one has*

$$\phi_!(\pi^*K \otimes \mathcal{P}_{W,\psi}^d) \xrightarrow{\sim} (W \otimes V_E^\gamma)^{(d)}[d + d_G](\frac{d + d_G}{2}) \quad \square$$

For a $\check{G}$ -local system E pick a $\check{G}$ -local system E^* such that $V_{E^*}^{\check{\lambda}} \xrightarrow{\sim} (V_E^{\check{\lambda}})^*$ for all $\check{\lambda} \in \check{\Lambda}^+$. Let K be a E -Hecke eigensheaf then $\mathbb{D}K$ is a E^* -Hecke eigensheaf. Assume that $\mathbb{D}K$ is generic normalized then from Corollary 1 we get an isomorphism

$$\phi_!(\pi^*(\mathbb{D}K) \otimes \mathcal{P}_{W,\psi}^d) \xrightarrow{\sim} (W \otimes V_{E^*}^\gamma)^{(d)}[d + d_G](\frac{d + d_G}{2})$$

By adjunction, it yields a nonzero map

$$\mathbb{D}K \rightarrow \pi_* \text{R}\mathcal{H}om(\mathcal{P}_{W,\psi}^d, \phi^!(W \otimes V_{E^*}^\gamma)^{(d)})[d + d_G](\frac{d + d_G}{2})$$

Dualizing, we see that this is equivalent to providing a nonzero map

$$\pi_!(\mathcal{P}_{W,\psi}^d \otimes \phi^*(W^* \otimes V_E^\gamma)^{(d)}) \rightarrow K[d_G - d](\frac{d_G - d}{2}) \quad (2)$$

Set $W = V_E^\gamma$, so we have a canonical map $\bar{\mathbb{Q}}_\ell \rightarrow (W^* \otimes V_E^\gamma)^{(d)}$ on $X^{(d)}$. Composing with (2) we get a morphism

$$\pi_!\mathcal{P}_{W,\psi}^d \rightarrow K[d_G - d](\frac{d_G - d}{2}) \quad (3)$$

Conjecture 1 (geometric Langlands). *Let E be a $\check{G}$ -local system on X . Assume that $W = V_E^\gamma$ is irreducible and satisfies the condition*

(A) *If E' is a $\check{G}$ -local system on X such that $V_{E'}^\gamma \xrightarrow{\sim} V_E^\gamma$ then $E' \xrightarrow{\sim} E$.*

Then there exists $N > 0$ and for each $d \geq N$ a nonempty open substack $U_d \subset \text{Bun}_G^d$ with the following property. There exists a E -Hecke eigensheaf K on Bun_G such that

- *both K and $\mathbb{D}K$ are generic normalized;*
- *for $d \geq N$ the complex $\pi_! \mathcal{P}_{W,\psi}^d|_{U_d}$ is placed in perverse degrees $\leq d - d_G$, and the map (3) induces an isomorphism*

$$\mathcal{H}^{d-d_G}(\pi_! \mathcal{P}_{W,\psi}^d) \xrightarrow{\sim} K|_{U_d}$$

on the top perverse cohomology sheaves;

- *K is an irreducible perverse sheaf over each Bun_G^d , which does not vanish over U_d .*

Remarks . i) The sheaf K from Conjecture 1 is unique up to an isomorphism if it exists.

ii) For any local system W on X we have $\pi_! \mathcal{P}_{W,\psi}^d \xrightarrow{\sim} \text{Av}_W^d(\pi_! \mathcal{P}_\psi^0)$ naturally.

2.8 INFORMAL MOTIVATION If the ground field k was finite then according to Langlands' spectral decomposition theorem ([13]), each function from $L^2(\text{Bun}_G(k))$ would be written as linear combination (more precisely, a direct integral) of Hecke eigenfunctions.

Conjecturally, some version of spectral decomposition should exist for the derived category $D(\text{Bun}_G)$ itself. We also have an analog of the scalar product of two objects $K_1, K_2 \in D(\text{Bun}_G)$, which is the cohomology $\text{R}\Gamma_c(\text{Bun}_G, K_1 \otimes \mathbb{D}(K_2))$ (we ignore here all convergence questions).

Let E be a $\check{G}$ -local system on X satisfying the assumptions of Conjecture 1. One may hope that to E is associated a E -Hecke eigensheaf K , which is unique in appropriate sense.

Since K is expected to be generic normalized, the "scalar product" of $\pi_! \mathcal{P}_\psi^0$ and K should equal "one". That is, K should appear in the spectral decomposition of $\pi_! \mathcal{P}_\psi^0$ with multiplicity one. By Proposition 3, the functor Av_W^d applied to $\pi_! \mathcal{P}_\psi^0$ with d large enough, will kill all the terms in the spectral decomposition of $\pi_! \mathcal{P}_\psi^0$ except K itself. So, roughly speaking, $\text{Av}_W^d(\pi_! \mathcal{P}_\psi^0)$ should equal K tensored by some constant complex.

2.9 STRATIFICATIONS For $\mu \in \Lambda^{\text{pos}}$ denote by X^μ the moduli scheme of Λ^{pos} -valued divisors of degree μ . If $\mu = \sum_{i \in \mathcal{I}} a_i \alpha_i$ then $X^\mu = \prod_{i \in \mathcal{I}} X^{(a_i)}$.

For $D \in X^{(d)}$ consider $\mathcal{F}'_T = \mathcal{F}_T(w_0(\gamma)D)$. The stack $\mathcal{Y}_d$ is the stack of pairs: $D \in X^{(d)}$ and a point $(\mathcal{F}_G, \kappa) \in \overline{\text{Bun}}_N^{\mathcal{F}'_T}$. Recall that $\overline{\text{Bun}}_N^{\mathcal{F}'_T}$ is stratified by locally closed substacks ${}_\mu \text{Bun}_N^{\mathcal{F}'_T}$ indexed by $\mu \in \Lambda^{\text{pos}}$. Namely, $(\mathcal{F}_G, \kappa)$ lies in ${}_\mu \text{Bun}_N^{\mathcal{F}'_T}$ iff there exists a divisor $D^{\text{pos}} \in X^\mu$ such that for all $\check{\lambda} \in \check{\Lambda}^+$ the meromorphic maps

$$\mathcal{L}_{\mathcal{F}'_T}^{\check{\lambda}}(\langle D^{\text{pos}}, \check{\lambda} \rangle) \rightarrow \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}$$

are regular everywhere and maximal. We have a projection

$${}_{\mu}\mathrm{Bun}_N^{\mathcal{F}'_T} \rightarrow X^{\mu}$$

whose fibre over D^{pos} is isomorphic to $\mathrm{Bun}_N^{\tilde{\mathcal{F}}_T}$, where $\tilde{\mathcal{F}}_T \xrightarrow{\sim} \mathcal{F}_T(w_0(\gamma)D + D^{pos})$.

Denote by ${}_{\mu}\mathcal{Y}_d \subset \mathcal{Y}_d$ the locally closed substack given by $(\mathcal{F}_G, \kappa) \in {}_{\mu}\mathrm{Bun}_N^{\mathcal{F}'_T}$. Let

$$\phi^{\mu} : {}_{\mu}\mathcal{Y}_d \rightarrow X^{(d)} \times X^{\mu}$$

be the projection. Let $X^{d,\mu} \subset X^{(d)} \times X^{\mu}$ be the closed subscheme given by the condition: $\gamma D + w_0(D^{pos})$ is dominant. This condition ensures that the maps $\tilde{\omega}_i : \mathcal{L}_{\tilde{\mathcal{F}}_T}^{\tilde{\alpha}_i} \rightarrow \Omega$ are regular for all $i \in \mathcal{I}$. Let ${}_{\mu}\mathcal{Y}_d^+ \subset {}_{\mu}\mathcal{Y}_d$ be the preimage of $X^{d,\mu}$ under ϕ^{μ} . So, we have the evaluation map

$$\mathrm{ev}_{\mu} : {}_{\mu}\mathcal{Y}_d^+ \rightarrow \mathbb{A}^1$$

Note that if $(D, D^{pos}) \in X^{d,\mu}$ then $\gamma D + w_0(D^{pos})$ is a $\Lambda_{G,S}^+$ -valued divisor of degree $d\gamma + w_0(\mu)$. In this way $X^{d,\mu}$ is the moduli scheme of $\Lambda_{G,S}^+$ -valued divisors on X of degree $d\gamma + w_0(\mu)$.

2.10 For any local system W on X let $\tilde{\mathcal{P}}_{W,\psi}^d$ be the complex obtained by replacing in the definition of $\mathcal{P}_{W,\psi}^d$ Laumon's sheaf by Springer's sheaf

$$\tilde{\mathcal{P}}_{W,\psi}^d = \mathfrak{q}_Y!(\mathcal{P}_{\psi}^0 \boxtimes \mathcal{S}pr_W^d)[-d_G](\frac{-d_G}{2})$$

Proposition 5. 1) For each $\mu \in \Lambda^{pos}$ the restriction of $\tilde{\mathcal{P}}_{W,\psi}^d$ to ${}_{\mu}\mathcal{Y}_d$ is supported by ${}_{\mu}\mathcal{Y}_d^+$ and is isomorphic to

$$\phi^{\mu*} \tilde{W}^{d,\mu} \otimes \mathrm{ev}_{\mu}^* \mathcal{L}_{\psi} \otimes (\bar{\mathbb{Q}}_{\ell}(\frac{1}{2})[1])^{\otimes d+d_N+\langle \gamma d - \mu, 2\tilde{\rho} \rangle}$$

for some sheaf $\tilde{W}^{d,\mu}$ on $X^{d,\mu}$. Here $\tilde{W}^{d,\mu}$ is placed in usual degree zero. Let $W^{d,\mu}$ denote the corresponding sheaves for $\mathcal{P}_{W,\psi}^d$.

2) If $(D = \sum d_k x_k, D^{pos} = \sum \mu_k x_k)$ is a k -point of $X^{d,\mu}$ then the fibre of $W^{d,\mu}$ at this point is the tensor product over all x_k

$$\otimes_k (\oplus_{\nu} \mathrm{Hom}(V^{d_k \gamma + w_0(\mu_k)}, (V^{\gamma})_{\nu}) \otimes (W_{x_k})_{\nu}),$$

the inside sum being taken over partitions ν of d_k of length $\leq r$.

The proof is found in Section 3.1. Proposition 5 together with Corolary 1 suggest the following conjecture.

Conjecture 2. Let $D = \sum \lambda_k x_k$ be a divisor on X with λ_k dominant coweights, let λ be the degree of D . Denote by $a : \mathrm{Bun}_N^{\tilde{\mathcal{F}}_T} \rightarrow \mathrm{Bun}_G$ the projection, where $\tilde{\mathcal{F}}_T = \mathcal{F}_T(w_0(D))$. Let $\mathrm{ev}_{\lambda} : \mathrm{Bun}_N^{\tilde{\mathcal{F}}_T} \rightarrow \mathbb{A}^1$ be the evaluation map given by the conductor data. Let E be a $\check{G}$ -local system on X , K be a generic normalized E -Hecke eigensheaf on Bun_G . Then

$$\mathrm{R}\Gamma_c(\mathrm{Bun}_N^{\tilde{\mathcal{F}}_T}, a^* K \otimes \mathrm{ev}_{\lambda}^* \mathcal{L}_{\psi}) \otimes (\bar{\mathbb{Q}}_{\ell}(\frac{1}{2})[1])^{d_N - d_G + \langle \lambda, 2\tilde{\rho} \rangle} \xrightarrow{\sim} \otimes_k (V_E^{\lambda_k})_{x_k}$$

2.11 Consider the diagram

$$\mathcal{Y}_d \times X \xleftarrow{\mathfrak{p}_Y \times \text{supp}} \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} \xrightarrow{\mathfrak{q}_Y} \mathcal{Y}_{d+1}$$

where $\mathfrak{p}_Y : \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} \rightarrow \mathcal{Y}_d$ is the projection. For any local system W on X define a natural map

$$(\mathfrak{p}_Y \times \text{supp})! \mathfrak{q}_Y^* \mathcal{P}_{W,\psi}^{d+1} \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes \langle \gamma, 2\check{\rho} \rangle} \rightarrow \mathcal{P}_{W,\psi}^d \boxtimes W \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1] \quad (4)$$

as follows. Consider the diagram

$$\begin{array}{ccc} \tilde{\mathcal{H}}_G^{+,d} \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} & \xrightarrow{\sim} & \tilde{\mathcal{H}}_G^{+,d+1} \\ \downarrow p \times \text{id} & & \downarrow p \\ \mathcal{H}_G^{+,d} \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} & \xrightarrow{\text{conv}} & \mathcal{H}_G^{+,d+1} \end{array}$$

Clearly, $\mathcal{L}_W^{d+1}$ is a direct summand of $\text{conv}!(\mathcal{L}_W^d \boxtimes \text{supp}^* W) \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes 1 + \langle \gamma, 2\check{\rho} \rangle}$. From the diagram

$$\begin{array}{ccc} \mathcal{Y}_0 \times_{\text{Bun}_G} \mathcal{H}_G^{+,d} \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} & \xrightarrow{\text{id} \times \text{conv}} & \mathcal{Y}_0 \times_{\text{Bun}_G} \mathcal{H}_G^{+,d+1} \\ \downarrow & & \downarrow \\ \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} & \xrightarrow{\mathfrak{q}_Y} & \mathcal{Y}_{d+1} \end{array}$$

we see that $\mathcal{P}_{W,\psi}^{d+1}$ is a direct summand of

$$\mathfrak{q}_Y^*(\mathcal{P}_{W,\psi}^d \boxtimes \text{supp}^* W) \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes 1 + \langle \gamma, 2\check{\rho} \rangle}$$

This yields a morphism

$$\mathfrak{q}_Y^* \mathcal{P}_{W,\psi}^{d+1} \rightarrow (\mathcal{P}_{W,\psi}^d \boxtimes \text{supp}^* W) \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes 1 + \langle \gamma, 2\check{\rho} \rangle}$$

Since $\mathfrak{p}_Y \times \text{supp} : \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} \rightarrow \mathcal{Y}_d \times X$ is smooth of relative dimension $\langle \gamma, 2\check{\rho} \rangle$, we get a map

$$\mathfrak{q}_Y^* \mathcal{P}_{W,\psi}^{d+1} \rightarrow (\mathfrak{p}_Y \times \text{supp})^!(\mathcal{P}_{W,\psi}^d \boxtimes W) \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes 1 - \langle \gamma, 2\check{\rho} \rangle}$$

and, by adjunction, the desired map (4).

Proposition 6. 1) *The complex*

$$(\mathfrak{p}_Y \times \text{supp})! \mathfrak{q}_Y^* \mathcal{P}_{W,\psi}^{d+1} \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes \langle \gamma, 2\check{\rho} \rangle} \quad (5)$$

is placed in perverse degrees ≤ 0 .

2) *For any $\mu \in \Lambda^{\text{pos}}$, the restriction of (5) to ${}_\mu \mathcal{Y}_d \times X$ is supported by ${}_\mu \mathcal{Y}_d^+ \times X$ and is isomorphic to the tensor product of*

$$\text{ev}_\mu^* \mathcal{L}_\psi \otimes (\bar{\mathbb{Q}}_\ell(\frac{1}{2})[1])^{\otimes d+1+d_N+\langle \gamma d - \mu, 2\check{\rho} \rangle}$$

with some sheaves $W_0^{d,\mu}$ coming from $X^{d,\mu} \times X$. Here $W_0^{d,\mu}$ is placed in usual degree zero.

The proof is given in Section 3.3

Remarks . i) One may show that for any $\mu \in \Lambda^{pos}$ the restriction of (4) to ${}_{\mu}\mathcal{Y}_d \times X$ comes from a morphism of sheaves $W_0^{d,\mu} \rightarrow W^{d,\mu} \boxtimes W$ on $X^{d,\mu} \times X$. The latter map is an isomorphism over the open substack of $X^{d,\mu} \times X$ classifying triples (D, D^{pos}, x) such that x does not appear in D . Therefore, (4) is an isomorphism over the locus of $(\mathcal{F}, \kappa, D, x) \in \mathcal{Y}_d \times X$ such that x does not appear in D .

ii) In the situation of Conjecture 1 we expect that the map (4) yields the Hecke property of K corresponding to the coweight γ . Moreover, it should also yield the Hecke properties corresponding to all $\lambda \in \Lambda_{G,S}^+$ (as it indeed happens for GL_n). Define the Hecke functor $\mathrm{H}_{\mathcal{Y}} : \mathrm{D}(\mathcal{Y}_{d+1}) \rightarrow \mathrm{D}(\mathcal{Y}_d \times X)$ by

$$\mathrm{H}_{\mathcal{Y}}(F) = (\mathfrak{p}_{\mathcal{Y}} \times \mathrm{supp})! \mathfrak{q}_{\mathcal{Y}}^* F \otimes \bar{\mathbb{Q}}_{\ell}(\frac{1}{2})[1]^{\otimes \langle \gamma, 2\check{\rho} \rangle}$$

Note that the cohomological shifts in the definition of the Hecke functor $\mathrm{H} : \mathrm{D}(\mathrm{Bun}_G^{d+1}) \rightarrow \mathrm{D}(\mathrm{Bun}_G^d \times X)$ corresponding to γ and $\mathrm{H}_{\mathcal{Y}}$ differ by one! So, the Hecke property of K can not be simply the push-forward of (4) with respect to $\pi : \mathcal{Y}_d \rightarrow \mathrm{Bun}_G$.

2.12 Let ω be a generator of the group of coweights orthogonal to all roots. Since the image of ω in $\pi_1(G)$ is not zero, we assume that this image equals $d_{\omega}\theta$ for some $d_{\omega} > 0$. Since $d_{\omega}\gamma - \omega$ is dominant, we have $d_{\omega}\gamma - \omega \in \Lambda^{pos}$ and $\omega \in \Lambda_{G,S}^+$. Consider the map $\mathfrak{q}^{\omega} : \mathrm{Bun}_G \times X \rightarrow \mathrm{Bun}_G$ sending $(\mathcal{F}_G, x)$ to $\mathcal{F}'_G$, where $\mathcal{F}'_G$ and $\mathcal{F}_G$ are identified over $X - x$ and

$$\mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}(\langle \omega, \check{\lambda} \rangle x) \xrightarrow{\sim} \mathcal{V}_{\mathcal{F}'_G}^{\check{\lambda}}$$

for each $\check{\lambda} \in \check{\Lambda}^+$. Let also

$$\mathfrak{q}_{\mathcal{Y}}^{\omega} : \mathcal{Y}_d \times X \rightarrow \mathcal{Y}_{d+d_{\omega}}$$

be the map sending $(\mathcal{F}_G, \kappa, x)$ to $(\mathcal{F}'_G, \kappa')$, where $\mathcal{F}'_G = \mathfrak{q}^{\omega}(\mathcal{F}_G, x)$ and $\kappa'^{\check{\lambda}}$ is the composition

$$\mathcal{L}_{\mathcal{F}_T}^{\check{\lambda}} \xrightarrow{\kappa^{\check{\lambda}}} \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}'_G}^{\check{\lambda}}$$

for all $\check{\lambda} \in \check{\Lambda}_S^+$. Let E be a $\check{G}$ -local system on X and set $W = V_E^{\gamma}$. Then there is a natural map

$$(\mathfrak{q}_{\mathcal{Y}}^{\omega})^* \mathcal{P}_{W,\psi}^{d+d_{\omega}} \otimes \bar{\mathbb{Q}}_{\ell}(\frac{1}{2})[1]^{\otimes 1-d_{\omega}} \rightarrow \mathcal{P}_{W,\psi}^d \boxtimes V_E^{\omega} \otimes \bar{\mathbb{Q}}_{\ell}(\frac{1}{2})[1] \quad (6)$$

This is not an isomorphism in general, and one may show that the LHS of (6) is placed in perverse degrees ≤ 0 .

2.13 WHITTAKER SHEAVES Let $\mathrm{K}(\mathcal{Y}_d)$ denote the Grothendieck ring of the triangulated category $\mathrm{D}(\mathcal{Y}_d)$.

To each $\check{G}$ -local system E on X and $d \geq 0$ we attach the Whittaker sheaf $\mathcal{W}_{E,\psi}^d \in \mathrm{K}(\mathcal{Y}_d)$ defined as follows.

Let $\mu \in \Lambda^{pos}$ be such that $d\gamma + w_0(\mu)$ is dominant. Let τ be a partition of $d\gamma + w_0(\mu)$, that is, a way to write $d\gamma + w_0(\mu) = \sum_i n_i \lambda_i$ with $\lambda_i \in \Lambda_{G,S}^+$ pairwise different and $n_i > 0$. Let ${}^\tau X \subset \prod_i X^{(n_i)}$ be the complement to all diagonals. We consider ${}^\tau X \subset X^{d,\mu}$ as the locally-closed subscheme classifying divisors $\sum_k \lambda_k x_k$ of degree $d\gamma + w_0(\mu)$ with $x_k \in X$ pairwise different.

Let ${}^\tau E$ denote the restriction of

$$\boxtimes_i (V_E^{\lambda_i})^{(n_i)}$$

under ${}^\tau X \hookrightarrow \prod_i X^{(n_i)}$. Let ${}^\tau \mathcal{Y} \subset {}_\mu \mathcal{Y}_d^+$ be the preimage of ${}^\tau X$ under $\phi^\mu : {}_\mu \mathcal{Y}_d^+ \rightarrow X^{d,\mu}$.

Definition 3. Set $\mathcal{W}_{E,\psi}^d \in K(\mathcal{Y}_d)$ to be the (unique) complex with the following properties. Its $*$ -restriction to each stratum ${}_\mu \mathcal{Y}_d$ is supported by ${}_\mu \mathcal{Y}_d^+$. If $d\gamma + w_0(\mu)$ is dominant then for any partition τ of $d\gamma + w_0(\mu)$ the $*$ -restriction of $\mathcal{W}_{E,\psi}^d$ to ${}^\tau \mathcal{Y}$ is

$$\phi^{\mu*}({}^\tau E) \otimes \text{ev}_\mu^* \mathcal{L}_\psi \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes d+d_N+\langle d\gamma-\mu, 2\check{\rho} \rangle}$$

The sheaf $\mathcal{W}_{E,\psi}^d$ should satisfy the Hecke property, in particular we suggest

Conjecture 3. Recall the diagram (cf. Sect. 2.11)

$$\mathcal{Y}_d \times X \xleftarrow{\mathfrak{p}\mathcal{Y} \times \text{supp}} \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} \xrightarrow{\mathfrak{q}\mathcal{Y}} \mathcal{Y}_{d+1}$$

There is a canonical isomorphism in the Grothendieck ring $K(\mathcal{Y}_d \times X)$

$$(\mathfrak{p}\mathcal{Y} \times \text{supp})! \mathfrak{q}\mathcal{Y}^* \mathcal{W}_{E,\psi}^{d+1} \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]^{\otimes \langle \gamma, 2\check{\rho} \rangle} \rightarrow \mathcal{W}_{E,\psi}^d \boxtimes V_E^\gamma \otimes \bar{\mathbb{Q}}_\ell(\frac{1}{2})[1]$$

2.13.1 We don't know if $\Lambda_{G,S}^+$ is a free semigroup in general, however this is the case for our examples $\text{GL}_n, \text{GSp}_{2n}, \text{GSpin}_{2n+1}$ (cf. appendix).

Assuming that $\Lambda_{G,S}^+$ is a free semigroup, we can describe $\mathcal{W}_{E,\psi}^d$ more precisely, namely "glue" the pieces on the strata ${}^\tau \mathcal{Y}$ to get a sheaf on ${}_\mu \mathcal{Y}_d^+$. To do so, we will glue the sheaves ${}^\tau E$ to get a constructible sheaf $AE^{d,\mu}$ on $X^{d,\mu}$ (here 'A' stands for 'automorphic').

Let $\lambda_1, \dots, \lambda_m$ be free generators of $\Lambda_{G,S}^+$ thus yielding $\Lambda_{G,S}^+ \xrightarrow{\sim} (\mathbb{Z}_+)^m$. Given $d \geq 0$ and $\mu \in \Lambda^{pos}$ with $d\gamma + w_0(\mu) = \sum_{i=1}^m a_i \lambda_i$ dominant, we get

$$X^{d,\mu} \xrightarrow{\sim} \prod_{i=1}^m X^{(a_i)}$$

Consider the sheaf

$$\boxtimes_{i=1}^m (V_E^{\lambda_i})^{(a_i)} \tag{7}$$

on $X^{d,\mu}$. Let $D = \sum_k \nu_k x_k$ be a k -point of $X^{d,\mu}$, where x_k are pairwise different and $\nu_k \in \Lambda_{G,S}^+$. Write $\nu_k = \sum_{i=1}^m a_{i,k} \lambda_i$ for each k . The fibre of (7) at D is

$$\otimes_{x_k} \otimes_{i=1}^m \text{Sym}^{a_{i,k}} (V_E^{\lambda_i})_{x_k}$$

There is (a unique up to a nonzero multiple) inclusion of $\check{G}$ -modules $V^{\nu_k} \hookrightarrow \bigotimes_{i=1}^m \text{Sym}^{a_{i,k}}(V^{\lambda_i})$. This yields a map

$$\bigotimes_{x_k} (V_E^{\nu_k})_{x_k} \hookrightarrow \bigotimes_{x_k} \bigotimes_{i=1}^m \text{Sym}^{a_{i,k}}(V_E^{\lambda_i})_{x_k} \quad (8)$$

The following is borrowed from [12].

Proposition 7. *Assume that $\Lambda_{G,S}^+$ is a free semigroup. There is a unique constructible subsheaf $AE^{d,\mu} \subset \bigboxtimes_{i=1}^m (V_E^{\lambda_i})^{(a_i)}$ whose fibre at any k -point $D = \sum_k \nu_k x_k$ of $X^{d,\mu}$ is the image of (8). $\square$*

3. SOME PROOFS

3.1 Recall that Gr_G is stratified by locally closed ind-subschemes S^μ indexed by all coweights $\mu \in \Lambda$. Informally, S^μ is the $N(\hat{K})$ -orbit of the point $\mu(t) \in \text{Gr}_G$, where $\hat{K} = k((t))$. We refer the reader to [3], Section 7.1 for the precise definition.

Recall the following notion from *loc.cit.*, section 7.1.4. Set $\hat{\mathcal{O}} = k[[t]]$. Let $\hat{\Omega}$ denote the completed module of relative differentials of $\hat{\mathcal{O}}$ over k (so, $\hat{\Omega}$ is a free $\hat{\mathcal{O}}$ -module generated by dt). Given a coweight $\eta \in \Lambda$ and isomorphisms

$$s_i : \hat{\mathcal{O}}(\langle \eta, \check{\alpha}_i \rangle) \xrightarrow{\sim} \hat{\Omega}$$

for each $i \in \mathcal{I}$, one defines an admissible character $\chi_\eta : N(\hat{K}) \rightarrow \mathbb{G}_a$ of conductor η as the sum $\chi_\eta = \sum_{i \in \mathcal{I}} \chi^i$, where $\chi^i : N(\hat{K}) \rightarrow \mathbb{G}_a$ is the composition

$$N(\hat{K}) \rightarrow N/[N, N](\hat{K}) \xrightarrow{u_i} \mathbb{G}_a(\hat{K}) \xrightarrow{s_i} \hat{\Omega}(\hat{K}) \xrightarrow{\text{Res}} \mathbb{G}_a$$

Here $u_i : N/[N, N] \rightarrow \mathbb{G}_a$ is the natural coordinate corresponding to the simple root $\check{\alpha}_i$. By (*loc.cit.*, Lemma 7.1.5), for $\nu \in \Lambda$ there exists a $(N(\hat{K}), \chi_\eta)$ -equivariant function $\chi_\eta^\nu : S^\nu \rightarrow \mathbb{A}^1$ if and only if $\nu + \eta \in \Lambda^+$. In the latter case this function is unique up to an additive constant.

Proof of Proposition 5 Let $(\sum d_k x_k, \mathcal{F}_G)$ be a k -point of $X^{(d)} \times \text{Bun}_G$. The fibre of $\text{supp} \times \mathfrak{q} : \mathcal{H}_G^{+,d} \rightarrow X^{(d)} \times \text{Bun}_G$ over this point identifies with

$$\prod_k \overline{\text{Gr}}_G^{d_k \gamma}$$

The restriction of Spr_W^d to this fibre is the tensor product of $(\bigotimes_k (W_{x_k}^{\otimes d_k}))[d + d_G](\frac{d+d_G}{2})$ with

$$\bigboxtimes_k \left(\bigoplus_{\nu \leq d_k \gamma} \mathcal{A}_\nu \otimes V_{\nu,k} \right)$$

where the inside sum is taken over dominant coweights $\nu \in \Lambda^+$ such that $\nu \leq d_k \gamma$, and $V_{\nu,k}$ are some vector spaces. Let

$${}^0 \mathfrak{q}_Y : \text{Bun}_N^{\mathcal{F}_T} \times_{\text{Bun}_G} \mathcal{H}_G^{+,d} \rightarrow \mathcal{Y}_d$$

denote the restriction of $\mathfrak{q}_Y$ to the open substack $\mathrm{Bun}_N^{\mathcal{F}_T} \times_{\mathrm{Bun}_G} \mathcal{H}_G^{+,d} \subset \mathcal{Y}_0 \times_{\mathrm{Bun}_G} \mathcal{H}_G^{+,d}$.

Fix a k -point $y \in \mathcal{Y}_d$, it is given by $(D = \sum d_k x_k, D^{pos} = \sum \mu_k x_k, \mathcal{F}_G, \kappa)$ such that x_k are pairwise different (and some of d_k may be zero). Let K_y denote the fibre at y of

$${}^0\mathfrak{q}_Y!(\mathrm{ev}^* \mathcal{L}_\psi \boxtimes \mathcal{S}pr_W^d)[d_N - d_G](\frac{d_N - d_G}{2})$$

The fibre of ${}^0\mathfrak{q}_Y$ over y identifies with

$$\prod_k S^{w_0(\gamma)d_k + \mu_k} \cap \overline{\mathrm{Gr}}_G^{d_k \gamma}$$

For each k set $\nu_k = -w_0(\gamma)d_k - \mu_k$. An equivariance argument (as in [3], Lemma 6.2.8) shows that K_y vanishes unless all ν_k are dominant.

By (Lemma 7.2.7(2), [3]) the restriction of the map

$$\mathrm{Bun}_N^{\mathcal{F}_T} \times_{\mathrm{Bun}_G} \mathcal{H}_G^{+,d} \rightarrow \mathrm{Bun}_N^{\mathcal{F}_T} \xrightarrow{\mathrm{ev}} \mathbb{A}^1$$

to $({}^0\mathfrak{q}_Y)^{-1}(y)$ becomes the sum over all k of

$$\chi_{\nu_k}^{-\nu_k} : S^{-\nu_k} \cap \overline{\mathrm{Gr}}^{d_k \gamma} \rightarrow \mathbb{A}^1$$

plus $\mathrm{ev}_\mu(\mathcal{F}_G, \kappa, D, D^{pos}) \in \mathbb{A}^1$. By (Theorem 1, [3]), for any $\nu \in \Lambda^+$ such that $\nu \leq d_k \gamma$ the complex

$$\mathrm{R}\Gamma_c(S^{-\nu_k} \cap \overline{\mathrm{Gr}}^{d_k \gamma}, \mathcal{A}_\nu \otimes (\chi_{\nu_k}^{-\nu_k})^* \mathcal{L}_\psi)$$

vanishes unless $\nu = -w_0(\nu_k)$. In the latter case, it is canonically $\bar{Q}_\ell[\langle \nu_k, 2\check{\rho} \rangle](\langle \nu_k, \check{\rho} \rangle)$.

The above equivariance argument shows also that the restriction of $\tilde{\mathcal{P}}_{W,\psi}^d$ to ${}_\mu\mathcal{Y}_d^+$, after tensoring by $\mathrm{ev}_\mu^* \mathcal{L}_{\psi^{-1}}$, descends with respect to the projection ${}_\mu\mathcal{Y}_d^+ \rightarrow X^{d,\mu}$. Combining this with Proposition 2, one finishes the proof of Proposition 5. $\square$

The above proof combined with (Proposition 3.2.6, [2]) also gives the following

Corollary 2. *Over the open substack ${}_0\mathcal{Y}_d \subset \mathcal{Y}_d$, the map ${}^0\mathfrak{q}_Y : \mathrm{Bun}_N^{\mathcal{F}_T} \times_{\mathrm{Bun}_G} \mathcal{H}_G^{+,d} \rightarrow \mathcal{Y}_d$ is an isomorphism.*

3.2 In this subsection we prove Proposition 4.

Given a pair $d \geq 0$, $\mu \in \Lambda^{pos}$ such that $d\gamma + w_0(\mu)$ is dominant, a *partition* τ of $d\gamma + w_0(\mu)$ is a presentation of $d\gamma + w_0(\mu)$ as a sum of nonzero elements from $\Lambda_{G,S}^+$, that is, $\sum_k (d_k \gamma + w_0(\mu_k)) = d\gamma + w_0(\mu)$ with $d_k \geq 0$, $\mu_k \in \Lambda^{pos}$ and $d_k \gamma + w_0(\mu_k)$ dominant for all k .

Given a partition τ of $d\gamma + w_0(\mu)$, consider the locally closed subscheme ${}^\tau X \subset X^{d,\mu}$, which is the moduli scheme of divisors $\sum_k (d_k \gamma + w_0(\mu_k))x_k$ with x_k pairwise different. Given τ , if k runs through the set consisting of m elements then $\dim {}^\tau X = m$. Clearly, the schemes ${}^\tau X$ form a stratification of $X^{d,\mu}$.

Let ${}^\tau\mathcal{Y} \subset {}_\mu\mathcal{Y}_d^+$ be the preimage of ${}^\tau X$ in ${}_\mu\mathcal{Y}_d^+$. Suppose that τ is of length m , that is, $\dim {}^\tau X = m$. Then from (Lemma 7.2.4, [3]) it follows that

$$\dim {}^\tau\mathcal{Y} = m + d_N + \langle \gamma d - \mu, 2\check{\rho} \rangle$$

In particular, we have $\dim \mathcal{Y}_d = d + d_N + \langle d\gamma, 2\check{\rho} \rangle$. Now from Proposition 5, we learn that the restriction of $\tilde{\mathcal{P}}_{W,\psi}^d$ to ${}^\tau\mathcal{Y}$ is placed in perverse degrees ≤ 0 . Moreover, the inequality is strict unless $\mu = 0$ and $m = d$. Since $\tilde{\mathcal{P}}_{W,\psi}^d$ is self-dual (up to replacing W by W^* and ψ by ψ^{-1}), our assertion follows. $\square$

Remarks . i) As a corollary, note that the restriction of $\mathcal{P}_{W,\psi}^d$ to ${}_0\mathcal{Y}_d$ identifies canonically with

$$\phi^{0*}W^{(d)} \otimes \text{ev}_0^* \mathcal{L}_\psi \otimes (\bar{\mathbb{Q}}_\ell(\frac{1}{2})[1])^{\otimes d+d_N+\langle \gamma d, 2\check{\rho} \rangle}$$

ii) For each pair $d \geq 0$, $\mu \in \Lambda^{pos}$ such that $d\gamma + w_0(\mu)$ is dominant, the projection $X^{d,\mu} \rightarrow X^{(d)}$ is a finite morphism. Let W and W' be any local systems on X . Then the complex

$$\phi_!(\mathcal{P}_{W,\psi}^d \otimes \mathcal{P}_{W',\psi^{-1}}^d) \quad (9)$$

is placed in usual cohomological degree $-2d$. This is seen by calculating this direct image with respect to the stratification of $\mathcal{Y}_d$ by ${}_\mu\mathcal{Y}_d$. For $G = \text{GL}_n$ the complex (9) is a Rankin-Selberg integral considered in [11].

3.3 In this subsection we prove Proposition 6.

Lemma 2. *Let $\lambda \in \Lambda^+$, $\mu \in \Lambda$. Let $w \in W$ be any element of the Weil group such that $\mu + w\lambda \in \Lambda^+$. The function $\chi_\mu^{w\lambda} : S^{w\lambda} \rightarrow \mathbb{A}^1$ is constant on $S^{w\lambda} \cap \overline{\text{Gr}}^\lambda$ if and only if $\mu \in \Lambda^+$.*

Proof This follows from the description of $S^{w\lambda} \cap \overline{\text{Gr}}^\lambda$ given in ([14], Lemma 5.2). $\square$

Proof of Proposition 6 The fibre of $\mathfrak{p} \times \text{supp} : \mathcal{H}_G^{+,1} \rightarrow \text{Bun}_G \times X$ over a k -point $(\mathcal{F}_G, x)$ is identified with $\text{Gr}_G^{-w_0(\gamma)}$. This fibre is stratified by the subschemes

$$S^{-w_0w(\gamma)} \cap \text{Gr}_G^{-w_0(\gamma)} \quad (10)$$

indexed by $w(\gamma) \in W\gamma$. By ([14], Lemma 5.2), (10) is an affine space of dimension $\langle \gamma + w(\gamma), \check{\rho} \rangle$.

Consider a k -point of ${}_\mu\mathcal{Y}_d$ given by $(\mathcal{F}_G, \kappa, D, D^{pos})$. Let $(\mathcal{F}_G, \mathcal{F}'_G, \beta, x \in X) \in \mathcal{H}_G^{+,1}$ correspond to a point of $S^{-w_0w(\gamma)} \cap \text{Gr}_G^{-w_0(\gamma)}$ for some $w(\gamma) \in W\gamma$. The image of this collection under the map

$$\mathfrak{q}\mathcal{Y} : \mathcal{Y}_d \times_{\text{Bun}_G} \mathcal{H}_G^{+,1} \rightarrow \mathcal{Y}_{d+1}$$

is the point of ${}_{\mu'}\mathcal{Y}_{d+1}$ given by

$$(\mathcal{F}'_G, \kappa', D + x, D^{pos} + w_0w(\gamma)x - w_0(\gamma)x)$$

with $\mu' = \mu + w_0 w(\gamma) - w_0(\gamma)$. If $\gamma D + w_0(D^{pos}) + w(\gamma)x$ is not dominant then, by Proposition 5, the stratum (10) does not contribute to the direct image (5).

Assume $\gamma D + w_0(D^{pos}) + w(\gamma)x$ dominant. Let $d_x \in \mathbb{Z}_+$ (resp, $\mu_x \in \Lambda^{pos}$) denote the multiplicity of x in D (resp., in D^{pos}). Then the restriction of $\text{ev}_{\mu'} \circ \mathbf{q}_{\mathcal{Y}}$ to the stratum (10) is $(N(\hat{K}), \chi_\nu)$ -equivariant for some admissible character $\chi_\nu : N(\hat{K}) \rightarrow \mathbb{G}_a$ of conductor $\nu = -w_0(\gamma)d_x - \mu_x$. If ν is dominant then this restriction is constant and equals $\text{ev}_\mu(\mathcal{F}_G, \kappa, D, D^{pos})$. If ν is not dominant then, by Lemma 2, the stratum (10) does not contribute to the direct image (5), because $\text{R}\Gamma_c(\mathbb{A}^1, \mathcal{L}_\psi) = 0$.

We conclude that the restriction of (5) to ${}_\mu \mathcal{Y}_d \times X$ vanishes outside the closed substack ${}_\mu \mathcal{Y}_d^+ \times X$, and is isomorphic to

$$F[d + 1 + d_N + \langle \gamma d - \mu, 2\check{\rho} \rangle]$$

for some sheaf F on ${}_\mu \mathcal{Y}_d^+ \times X$ placed in usual degree zero. An equivariance argument (as in the proof of Proposition 5) assures that $F \otimes \text{ev}_\mu^* \mathcal{L}_{\psi^{-1}}$ descends with respect to the projection ${}_\mu \mathcal{Y}_d^+ \times X \rightarrow X^{d, \mu} \times X$.

Since $\dim({}_\mu \mathcal{Y}_d^+) \leq d + d_N + \langle \gamma d - \mu, 2\check{\rho} \rangle$ for any $\mu \in \Lambda^{pos}$, the complex (5) is placed in perverse degrees ≤ 0 . Proposition 6 is proved. $\square$

4. ADDITIONAL STRUCTURE ON LEVI SUBGROUPS

4.1 Throughout this section, we fix a standard parabolic subgroup $P \subset G$ corresponding to a subset $\mathcal{I}_M \subset \mathcal{I}$. Let $\Lambda_{G,P}$ be the quotient of Λ by the $\mathbb{Z}$ -span of $\alpha_i, i \in \mathcal{I}_M$. Let

$$\check{\Lambda}_{G,P} = \{\check{\lambda} \in \check{\Lambda} \mid \langle \alpha_i, \check{\lambda} \rangle = 0 \text{ for } i \in \mathcal{I}_M\}$$

be the dual lattice. Let $U(P)$ be the unipotent radical of P and $M = P/U(P)$. Write $\Lambda_M^+ \subset \Lambda$ for the semigroup of dominant coweights for M , and w_0^M for the longest element of the Weil group W_M of M . The irreducible $\check{M}$ -module of h.w. λ is denoted U^λ . Fix a section $M \rightarrow P$ of the projection $P \rightarrow M$.

Definition 4. Set $\Lambda_{M,S}^+ = \{\lambda \in \Lambda_M^+ \mid \text{there exists } w \in W \text{ such that } w\lambda \in \Lambda_{G,S}^+\}$. Let $\pi_1^+(M) \subset \pi_1(M) = \Lambda_{G,P}$ denote the image of the projection $\Lambda_{M,S}^+ \rightarrow \Lambda_{G,P}$. Set

$$\check{\Lambda}_{M,S}^+ = \{\check{\lambda} \in \check{\Lambda}_M^+ \mid \text{there exists } w \in W \text{ such that } w\check{\lambda} \in \check{\Lambda}_S^+\}$$

The following identities are straightforward.

Lemma 3. *We have*

$$\Lambda_{M,S}^+ = \{\lambda \in \Lambda_M^+ \mid \langle w(\lambda), \check{\lambda} \rangle \geq 0 \text{ for all } w \in W, \check{\lambda} \in \check{\Lambda}_S^+\}$$

$$\Lambda_{M,S}^+ = \{\lambda \in \Lambda_M^+ \mid \langle w_0^M(\lambda), \check{\lambda} \rangle \geq 0 \text{ for all } \check{\lambda} \in \check{\Lambda}_{M,S}^+\}$$

$$\check{\Lambda}_{M,S}^+ = \{\check{\lambda} \in \check{\Lambda}_M^+ \mid \langle w(\lambda), \check{\lambda} \rangle \geq 0 \text{ for all } w \in W, \lambda \in \Lambda_{G,S}^+\}$$

$$\check{\Lambda}_{M,S}^+ = \{\check{\lambda} \in \check{\Lambda}_M^+ \mid \langle w_0^M(\lambda), \check{\lambda} \rangle \geq 0 \text{ for all } \lambda \in \Lambda_{M,S}^+\} \quad \square$$

Definition 5. Let $\text{Gr}_M^+ \subset \text{Gr}_M$ be the closed subscheme given by the condition: $(\mathcal{F}_M, \beta) \in \text{Gr}_M^+$ iff for each $\check{\lambda} \in \check{\Lambda}_{M,S}^+$ the map

$$\beta^{\check{\lambda}} : U_{\mathcal{F}_M}^{\check{\lambda}} \rightarrow U_{\mathcal{F}_M^0}^{\check{\lambda}}$$

is regular on $\mathcal{D}$. Here $U^{\check{\lambda}}$ stands for the irreducible quotient of the Weil module $\mathcal{U}^{\check{\lambda}}$ for M .

Clearly, Gr_M^+ is a $M(\hat{\mathcal{O}})$ -invariant subscheme of Gr_M . For $\nu \in \Lambda_M^+$ one has the closed subscheme $\overline{\text{Gr}}_M^\nu \subset \text{Gr}_M$ (cf. [2], sect. 3.2). For $\nu \in \Lambda_M^+$ we have $\overline{\text{Gr}}_M^\nu \subset \text{Gr}_M^+$ iff $\nu \in \Lambda_{M,S}^+$.

Recall that for $\mu \in \pi_1(M)$ the connected component Gr_M^μ of Gr_M classifies $(\mathcal{F}_M, \beta) \in \text{Gr}_M$ such that for $\check{\lambda} \in \check{\Lambda}_{G,P}$ we have

$$\mathcal{V}_{\mathcal{F}_M^0}^{\check{\lambda}}(-\langle \nu, \check{\lambda} \rangle) \xrightarrow{\sim} \mathcal{V}_{\mathcal{F}_M}^{\check{\lambda}}$$

For $\mu \in \pi_1(M)$ set $\text{Gr}_M^{+,\mu} = \text{Gr}_M^+ \cap \text{Gr}_M^\mu$. So, $\text{Gr}_M^{+,\mu}$ is nonempty iff $\mu \in \pi_1^+(M)$. It may be shown that $\text{Gr}_M^{+,\mu}$ is connected for each $\mu \in \pi_1^+(M)$.

Recall the following definition from ([2], sect. 4.3.1). For $\mu \in \Lambda_{G,P}$ let $S_P^\mu \subset \text{Gr}_G$ denote the locally closed subscheme classifying $(\mathcal{F}_G, \beta : \mathcal{F}_G^0|_{\mathcal{D}^*} \xrightarrow{\sim} \mathcal{F}_G|_{\mathcal{D}^*}) \in \text{Gr}_G$ such that the composition

$$\mathcal{L}_{\mathcal{F}_T^0}^{\check{\lambda}}(-\langle \mu, \check{\lambda} \rangle) \rightarrow \mathcal{L}_{\mathcal{F}_T^0}^{\check{\lambda}} \rightarrow \mathcal{V}_{\mathcal{F}_G^0}^{\check{\lambda}} \xrightarrow{\beta} \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}$$

has neither pole nor zero over $\mathcal{D}$ for every $\check{\lambda} \in \check{\Lambda}_{G,P} \cap \check{\Lambda}^+$.

For each $\nu \in \pi_1(G)$ the component Gr_G^ν is stratified by S_P^μ indexed by those $\mu \in \Lambda_{G,P}$ whose image in $\pi_1(G)$ is ν . Moreover, we have a natural map $t_S^\mu : S_P^\mu \rightarrow \text{Gr}_M^\mu$.

Lemma 4. For each $\mu \in \Lambda_{G,P}$ the map $t_S^\mu : \text{Gr}_G^+ \cap S_P^\mu \rightarrow \text{Gr}_M^\mu$ factors through $\text{Gr}_M^{+,\mu} \hookrightarrow \text{Gr}_M^\mu$, and the induced map $t_S^{+,\mu} : \text{Gr}_G^+ \cap S_P^\mu \rightarrow \text{Gr}_M^{+,\mu}$ is surjective.

Proof Let $(\mathcal{F}_G, \beta : \mathcal{F}_G^0|_{\mathcal{D}^*} \rightarrow \mathcal{F}_G|_{\mathcal{D}^*}) \in \text{Gr}_G^+ \cap S_P^\mu$. So, for any $\check{\lambda} \in \check{\Lambda}_{G,P} \cap \check{\Lambda}^+$

$$\mathcal{L}_{\mathcal{F}_T^0}^{\check{\lambda}}(-\langle \mu, \check{\lambda} \rangle) \hookrightarrow \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}$$

is a subbundle. There is unique $(\mathcal{F}_P, \beta : \mathcal{F}_P^0|_{\mathcal{D}^*} \rightarrow \mathcal{F}_P|_{\mathcal{D}^*}) \in \text{Gr}_P$ that induces $(\mathcal{F}_G, \beta)$, and t_S^μ sends $(\mathcal{F}_G, \beta)$ to $\mathcal{F}_M = \mathcal{F}_P \times_P M$. Since for any $\check{\lambda} \in \check{\Lambda}_S^+$ the maps

$$\beta^{\check{\lambda}} : \mathcal{V}_{\mathcal{F}_P}^{\check{\lambda}} \hookrightarrow \mathcal{V}_{\mathcal{F}_P^0}^{\check{\lambda}}$$

are regular, the first assertion is reduced to the next sublemma.

Sublemma 1. Let $\check{\nu} \in \check{\Lambda}_{M,S}^+$. So, there exists $w \in W$ with $w\check{\nu} \in \check{\Lambda}_S^+$. Let $\text{Res}_G^P V^{w\check{\nu}}$ denote $V^{w\check{\nu}}$ viewed as a P -module. There exists a subquotient V' of $\text{Res}_G^P V^{\check{\lambda}}$ on which $U(P)$ acts trivially and such that

$$\text{Hom}_M(U^{\check{\nu}}, V') \neq 0 \quad \square$$

Let $\nu \in \Lambda_{M,S}^+$ be such that $\text{Gr}_M^\nu \subset \text{Gr}_M^{+,\mu}$. Recall the notation $\hat{\mathcal{O}} = k[[t]]$. Since $t_S^{+,\mu}$ is $M(\hat{\mathcal{O}})$ -invariant, it suffices to show that $\nu(t) \in \text{Gr}_M^{+,\mu}$ lies in the image of $t_S^{+,\mu}$. We know that there exists $w \in W$ with $w\nu \in \Lambda_{G,S}^+$. Therefore, $\nu(t)G(\hat{\mathcal{O}})$ defines a point of $\text{Gr}_G^+ \cap S_P^\mu$ which is sent by $t_S^{+,\mu}$ to $\nu(t) \in \text{Gr}_M^{+,\mu}$.

□(Lemma 4)

Note as a consequence that for each $\nu \in \pi_1^+(G)$ the scheme $\text{Gr}_G^{+,\nu}$ is stratified by locally closed subschemes $\text{Gr}_G^{+,\nu} \cap S_P^\mu$ indexed by those $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is ν .

For $d \geq 0$ write $\Lambda_{M,S}^{+,d\theta}$ for the preimage of $d\theta$ under $\Lambda_{M,S}^+ \rightarrow \pi_1^+(G)$.

Lemma 5. *For any $\lambda \in \Lambda_{M,S}^{+,d\theta}$ there exist $\lambda_1, \dots, \lambda_d \in \Lambda_{M,S}^{+,\theta}$ such that $\lambda \leq_M \lambda_1 + \dots + \lambda_d$.*

Proof Pick any k -point $(\mathcal{F}_M, \beta)$ of Gr_M^λ . Let μ be the image of λ in $\pi_1^+(M)$. Pick any k -point $(\mathcal{F}_G, \beta)$ of $\text{Gr}_G^{+,d} \cap S_P^\mu$ whose $t_S^{+,\mu}$ -image is $(\mathcal{F}_M, \beta)$. Let $\text{Gr}_G^\gamma \tilde{\times} \dots \tilde{\times} \text{Gr}_G^\gamma$ be the scheme classifying collections

$$(\mathcal{F}_G^1, \dots, \mathcal{F}_G^{d+1} = \mathcal{F}_G^0, \beta_i), \quad (11)$$

where $\mathcal{F}_G^i$ is a G -bundle on $\mathcal{D}$, and $\beta^i : \mathcal{F}_G^i|_{\mathcal{D}^*} \xrightarrow{\sim} \mathcal{F}_G^{i+1}|_{\mathcal{D}^*}$ is an isomorphism such that $\mathcal{F}_G^i$ is in the position γ with respect to $\mathcal{F}_G^{i+1}$ for $i = 1, \dots, d$.

Pick a k -point (11) whose image under the convolution map

$$\text{Gr}_G^\gamma \tilde{\times} \dots \tilde{\times} \text{Gr}_G^\gamma \rightarrow \overline{\text{Gr}}_G^{d\gamma},$$

is $(\mathcal{F}_G, \beta)$. There exist a unique collection

$$(\mathcal{F}_P^1, \dots, \mathcal{F}_P^d, \mathcal{F}_P^{d+1} = \mathcal{F}_P^0, \beta_i),$$

where $\mathcal{F}_P^i$ is a P -torsor on $\mathcal{D}$ and $\beta_i : \mathcal{F}_P^i|_{X-x_i} \xrightarrow{\sim} \mathcal{F}_P^{i+1}|_{X-x_i}$, that induces (11) by extension of scalars from P to G . Extending the scalars from P to M , one gets a collection

$$(\mathcal{F}_M^1, \dots, \mathcal{F}_M^d, \mathcal{F}_M^{d+1} = \mathcal{F}_M^0, \beta_i)$$

where $\mathcal{F}_M^1 = \mathcal{F}_M$. For $i = 1, \dots, d$ let $\lambda_i \in \Lambda_M^+$ be such that $\mathcal{F}_M^i$ is in the position λ_i with respect to $\mathcal{F}_M^{i+1}$. By Lemma 4, $\lambda_i \in \Lambda_{M,S}^{+,\theta}$ for all i . Let

$$\text{Gr}_M^{\lambda_1} \tilde{\times} \dots \tilde{\times} \text{Gr}_M^{\lambda_d}$$

be the scheme classifying $(\mathcal{F}_M^1, \dots, \mathcal{F}_M^{d+1} = \mathcal{F}_M^0, \beta^i)$, where $\mathcal{F}_M^i$ is a M -torsor on $\mathcal{D}$ and $\beta^i : \mathcal{F}_M^i|_{\mathcal{D}^*} \xrightarrow{\sim} \mathcal{F}_M^{i+1}|_{\mathcal{D}^*}$ is an isomorphism such that $\mathcal{F}_M^i$ is in the position λ_i with respect to $\mathcal{F}_M^{i+1}$ for $i = 1, \dots, d$. We learn that $(\mathcal{F}_M, \beta)$ lies in the image of the convolution map

$$\text{Gr}_M^{\lambda_1} \tilde{\times} \dots \tilde{\times} \text{Gr}_M^{\lambda_d} \rightarrow \text{Gr}_M^{+,\mu}$$

But the image of the latter map is contained in $\overline{\text{Gr}}_M^{\lambda_1 + \dots + \lambda_d}$, so $\text{Gr}_M^\lambda \subset \overline{\text{Gr}}_M^{\lambda_1 + \dots + \lambda_d}$. □

Denote by $\pi_1^\theta(M)$ the image of $\Lambda_{M,S}^{+,\theta} \rightarrow \pi_1^+(M)$. By the above lemma, $\pi_1^\theta(M)$ generates $\pi_1^+(M)$ as a semigroup. Since V^γ is faithful, $\pi_1^\theta(M)$ generates $\pi_1(M)$ as a group.

Lemma 6. 1) Each $\lambda \in \Lambda_{M,S}^{+,\theta}$ is a minuscule dominant coweight for M .

2) The natural map $\Lambda_{M,S}^{+,\theta} \rightarrow \pi_1^\theta(M)$ is bijective.

Proof 1) For $\lambda \in \Lambda_{M,S}^{+,\theta}$ we have $\text{Hom}_M(U^\lambda, V^\gamma) \neq 0$. Let $\lambda' \in \Lambda_M^+$ and $\lambda' \leq_M \lambda$. Then λ' is a weight of V^γ , so that $\lambda' = w\gamma$ for some $w \in W$, and $\lambda = w\gamma + \alpha$, where α is a sum of positive coroots for M . However, if $\alpha \neq 0$ then the length of λ is strictly bigger than the length of γ , which implies that λ is not a weight of V^γ . This contradiction shows that $\lambda' = \lambda$.

2) follows from 1). $\square$

4.2 For $d \geq 0$ consider the stack

$$\mathcal{H}_G^{+,d} \times_{\text{Bun}_G} \text{Bun}_P, \quad (12)$$

where we used the projection $\mathfrak{q} : \mathcal{H}_G^{+,d} \rightarrow \text{Bun}_G$ in the fibred product.

For $\mu \in \Lambda_{G,P}$ let $\mathcal{H}_P^{+,\mu}$ be the locally closed substack of (12) classifying

$$(D \in X^{(d)}, \mathcal{F}_G, \mathcal{F}'_P, \beta : \mathcal{F}_G|_{X-D} \xrightarrow{\sim} \mathcal{F}'_G|_{X-D}, \mathcal{F}'_P \times_P G \xrightarrow{\sim} \mathcal{F}'_G)$$

for which there exists a $\Lambda_{G,P}$ -valued divisor D^μ on X of degree μ with the property: for all $\check{\lambda} \in \check{\Lambda}_{G,P} \cap \check{\Lambda}^+$ the meromorphic maps

$$\mathcal{L}_{\mathcal{F}'_{M/[M,M]}}^{\check{\lambda}} \rightarrow \mathcal{V}_{\mathcal{F}'_G}^{\check{\lambda}} \rightarrow \mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}$$

realise $\mathcal{L}_{\mathcal{F}'_{M/[M,M]}}^{\check{\lambda}}$ as a subbundle of $\mathcal{V}_{\mathcal{F}_G}^{\check{\lambda}}(\langle D^\mu, \check{\lambda} \rangle)$. Here we have denoted $\mathcal{F}'_M = \mathcal{F}'_P \times_P M$.

Clearly, we have $\langle D^\mu, \check{\omega}_0 \rangle = D$ and $\langle D^\mu, \check{\omega}_i \rangle \geq 0$ for $i \in \mathcal{I} - \mathcal{I}_M$. Let $D_i = \langle D^\mu, \check{\omega}_i \rangle$ for $i \in \mathcal{I} - \mathcal{I}_M$ then we have an equality of $\pi_1(M)$ -valued divisors on X

$$D^\mu = w_0(\gamma)D + \sum_{i \in \mathcal{I} - \mathcal{I}_M} D_i \alpha_i$$

By Lemma 4, $\mathcal{H}_P^{+,\mu}$ is non empty iff $\mu \in \pi_1^+(M)$ and actually D^μ is a $\pi_1^+(M)$ -valued divisor on X . So, for each $d \geq 0$ the stack (12) is stratified by locally closed substacks $\mathcal{H}_P^{+,\mu}$ indexed by those $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is $d\theta$.

For $\mu \in \pi_1^+(M)$ let $d = \langle \mu, \check{\omega}_0 \rangle$ and $d_i = \langle \mu, \check{\omega}_i \rangle$ for $i \in \mathcal{I} - \mathcal{I}_M$ and let X_M^μ denote the scheme image of the projection

$$\mathcal{H}_P^{+,\mu} \rightarrow X^{(d)} \times \prod_{i \in \mathcal{I} - \mathcal{I}_M} X^{(d_i)}$$

We will think of X_M^μ as the moduli scheme of $\pi_1^+(M)$ -valued divisors on X of degree μ . As we will see, X_M^μ need not be irreducible. For $\mu \in \pi_1^+(M)$ we have a commutative diagram

$$\begin{array}{ccc} \mathcal{H}_P^{+,\mu} & \hookrightarrow & \mathcal{H}_G^{+,d} \times_{\text{Bun}_G} \text{Bun}_P \\ \downarrow \text{supp}_P & & \downarrow \\ X_M^\mu & \xrightarrow{s_M} & X^{(d)}, \end{array}$$

where we have denoted by supp_P and s_M the natural projections.

Definition 6. For $\mu \in \pi_1^+(M)$ let $\mathcal{H}_M^{+,\mu}$ be the stack of collections $(\mathcal{F}_M, \mathcal{F}'_M, D^\mu \in X_M^\mu, \beta)$, where $\mathcal{F}_M, \mathcal{F}'_M \in \text{Bun}_M$ and

$$\beta : \mathcal{F}_M|_{X-D} \xrightarrow{\sim} \mathcal{F}'_M|_{X-D}$$

is an isomorphism of M -torsors with $D = s_M(D^\mu)$ such that for each $\check{\lambda} \in \check{\Lambda}_{M,S}^+$ the map

$$\beta^{\check{\lambda}} : U_{\mathcal{F}_M}^{\check{\lambda}} \hookrightarrow U_{\mathcal{F}'_M}^{\check{\lambda}}$$

extends to an inclusion of coherent sheaves on X , and

$$\mathcal{L}_{\mathcal{F}_M/[M,M]}^{\check{\lambda}}(\langle D^\mu, \check{\lambda} \rangle) \xrightarrow{\sim} \mathcal{L}_{\mathcal{F}'_M/[M,M]}^{\check{\lambda}}$$

for each $\check{\lambda} \in \check{\Lambda}_{G,P}$.

We have a diagram

$$\begin{array}{ccccc} \text{Bun}_M & \xleftarrow{\mathfrak{p}_M} & \mathcal{H}_M^{+,\mu} & \xrightarrow{\mathfrak{q}_M} & \text{Bun}_M \\ & & \downarrow \text{supp}_M & & \\ & & X_M^\mu & & \end{array}$$

where $\mathfrak{p}_M$ (resp., $\mathfrak{q}_M$) sends a point of $\mathcal{H}_M^{+,\mu}$ to $\mathcal{F}_M$ (resp., to $\mathcal{F}'_M$), and supp_M stands for the projection. If $(\mathcal{F}'_M, D^\mu = \sum_k \mu_k x_k)$ is a k -point of $\text{Bun}_M \times X_M^\mu$ then the fibre of

$$\mathfrak{q}_M \times \text{supp}_M : \mathcal{H}_M^{+,\mu} \rightarrow \text{Bun}_M \times X_M^\mu$$

over it identifies with $\prod_k \text{Gr}_M^{+,\mu_k}$.

4.3 Given $\mu \in \pi_1^+(M)$, we will denote by $\mathfrak{A}(\mu)$ the elements of the set of decompositions of μ as a sum of non-zero elements of $\pi_1^+(M)$. More precisely, $\mathfrak{A}(\mu)$ is a way to write $\mu = \sum_k n_k \mu_k$, where all $n_k > 0$ and $\mu_k \in \pi_1^+(M) - \{0\}$ are pairwise distinct.

For $\mathfrak{A}(\mu)$ we set $X^{\mathfrak{A}(\mu)} = \prod_k X^{(n_k)}$. We have a natural map $s^{\mathfrak{A}(\mu)} : X^{\mathfrak{A}(\mu)} \rightarrow X_M^\mu$ sending $\{D_k\}$ to $\sum_k \mu_k D_k$. Let $\check{X}^{\mathfrak{A}(\mu)} \subset X^{\mathfrak{A}(\mu)}$ be the complement to all diagonals. The composition

$$\check{X}^{\mathfrak{A}(\mu)} \hookrightarrow X^{\mathfrak{A}(\mu)} \rightarrow X_M^\mu$$

is a locally closed embedding, and in this way X_M^μ is stratified by subschemes $\check{X}^{\mathfrak{A}(\mu)}$.

We say that $\mathfrak{A}(\mu)$ is in general position if $\sum_k n_k = d$. Write ${}^{rss}X_M^\mu$ for the preimage of ${}^{rss}X^{(d)}$ under $s_M : X_M^\mu \rightarrow X^{(d)}$. The connected components of ${}^{rss}X_M^\mu$ are exactly $\check{X}^{\mathfrak{A}(\mu)}$ indexed by $\mathfrak{A}(\mu)$ in general position.

Definition 7. Given a local system W on X , for each $\mu \in \pi_1^+(M)$ define Laumon's sheaf $\mathcal{L}_W^\mu$ on $\mathcal{H}_M^{+,\mu}$ as follows. Recall the diagram

$$\mathcal{H}_M^{+,\mu} \xrightarrow{\text{supp}_M} X_M^\mu \xrightarrow{s_M} X^{(d)}$$

Let ${}^{rss}\mathcal{H}_M^{+, \mu}$ be the preimage of ${}^{rss}X^{(d)}$ under $s_M \circ \text{supp}_M$. The stack ${}^{rss}\mathcal{H}_M^{+, \mu}$ is smooth and over it we let

$$\mathcal{L}_W^\mu = \text{supp}_M^* s_M^* W^{(d)}[a](\frac{a}{2}),$$

where a denotes the dimension of the corresponding connected component of ${}^{rss}\mathcal{H}_M^{+, \mu}$. Then we extend this perverse sheaf by Goresky-MacPherson to $\mathcal{H}_M^{+, \mu}$.

By definition, $\mathbb{D}(\mathcal{L}_W^\mu) \xrightarrow{\sim} \mathcal{L}_{W^*}^\mu$, and $\mathcal{L}_W^\mu \xrightarrow{\sim} \oplus \mathcal{L}_W^{\mathfrak{A}(\mu)}$ is a direct sum of perverse sheaves indexed by $\mathfrak{A}(\mu)$ in general position. Set $d_M = \dim \text{Bun}_M$.

For $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is $d\theta$ consider the diagram

$$\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P \xleftarrow{q_M} \mathcal{H}_P^{+, \mu} \xrightarrow{f_M} \mathcal{H}_G^{+, d}$$

where we used $q_M : \mathcal{H}_M^{+, \mu} \rightarrow \text{Bun}_M$ in the fibred product, f_M is the natural map, and q_M sends $(\mathcal{F}_P, \mathcal{F}'_P, D^\mu, \beta)$ to $(\mathcal{F}_M, \mathcal{F}'_P, D^\mu, \mathcal{F}'_P \times_P M \xrightarrow{\sim} \mathcal{F}'_M, \beta)$. We also have

$$\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P \xleftarrow{p_M} \mathcal{H}_P^{+, \mu} \xrightarrow{f_M} \mathcal{H}_G^{+, d},$$

where now we used $p_M : \mathcal{H}_M^{+, \mu} \rightarrow \text{Bun}_M$ in the fibred product, and p_M sends $(\mathcal{F}_P, \mathcal{F}'_P, D^\mu, \beta)$ to $(\mathcal{F}'_M, \mathcal{F}_P, D^\mu, \mathcal{F}_P \times_P M \xrightarrow{\sim} \mathcal{F}_M, \beta)$.

Here is a generalization of Laumon's theorem ([10], Theorem 4.1).

Proposition 8. *Let W be a local system on X . Let $\mu \in \pi_1^+(M)$ with image $d\theta$ in $\pi_1^+(G)$. The complex $q_{M!} \mathcal{L}_W^d$ is canonically isomorphic to the inverse image of*

$$\mathcal{L}_W^\mu \otimes \bar{\mathbb{Q}}_\ell[1](\frac{1}{2})^{\otimes d_G - d_M + \langle \mu, 2\check{\rho}_M - 2\check{\rho} \rangle}$$

under the projection $\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P \rightarrow \mathcal{H}_M^{+, \mu}$. (We have used the fact that $\check{\rho}_M - \check{\rho} \in \check{\Lambda}_{G,P}$). The complex $p_{M!} \mathcal{L}_W^d$ is canonically isomorphic to the inverse image of

$$\mathcal{L}_W^\mu \otimes \bar{\mathbb{Q}}_\ell[1](\frac{1}{2})^{\otimes d_G - d_M - \langle \mu, 2\check{\rho}_M - 2\check{\rho} \rangle}$$

under the projection $\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P \rightarrow \mathcal{H}_M^{+, \mu}$.

The proof is given in Sections 4.4-4.5.

4.4 Let $J = \{i \in \mathcal{I} \mid \langle \gamma, \check{\alpha}_i \rangle = 0\}$. Let $W_J \subset W$ be the subgroup generated by the reflection corresponding to $i \in J$. Using Bruhat decomposition, one checks that the map $W/W_J \rightarrow W\gamma$ sending w to $w\gamma$ is a bijection.

Fix a section $T \rightarrow B$. Let P_γ denote the parabolic of G generated by T and $U_{\check{\alpha}}$ for all roots $\check{\alpha}$ such that $\langle \gamma, \check{\alpha} \rangle \leq 0$. So, P_γ contains the opposite Borel. We have a bijection $\Lambda_{M,S}^{+, \theta} \xrightarrow{\sim} W_M \backslash W/W_J$ sending $w\gamma \in \Lambda_{M,S}^{+, \theta}$ to the coset $W_M w W_J$.

The map $G/P_\gamma \rightarrow \text{Gr}_G^\gamma$ sending $g \in G(k) \subset G(\hat{\mathcal{O}})$ to $g\gamma(t)G(\hat{\mathcal{O}})$ is an isomorphism. The scheme Gr_G^γ is stratified by $\text{Gr}_G^\gamma \cap S_P^\lambda$ indexed by $\lambda \in \Lambda_{M,S}^{+,\theta}$. The above isomorphism transforms this stratification into the stratification of G/P_γ by P -orbits. We have a disjoint decomposition

$$G = \bigsqcup_{w \in W_M \setminus W/W_J} PwP_\gamma$$

So, we have

$$\text{Gr}_G^\gamma \cap S_P^\lambda \xrightarrow{\sim} PwP_\gamma/P_\gamma \xrightarrow{\sim} P/P \cap wP_\gamma w^{-1}$$

Similarly, for $\lambda \in \Lambda_{M,S}^{+,\theta}$ let $P_\lambda(M)$ be the parabolic of M generated by T and $U_{\check{\alpha}}$, where $\check{\alpha}$ runs through those roots of M for which $\langle \lambda, \check{\alpha} \rangle \leq 0$. Then the map $M/P_\lambda(M) \rightarrow \text{Gr}_M^\lambda$ sending $m \in M(k)$ to $m\lambda(t)M(\hat{\mathcal{O}})$ is an isomorphism. So, the map

$$\text{Gr}_G^\gamma \cap S_P^{w\gamma} \rightarrow \text{Gr}_M^{w\gamma} \quad (13)$$

is nothing else but the map $P/P \cap wP_\gamma w^{-1} \rightarrow M/P_{w\gamma}(M)$ sending p to $p \bmod M$. The correctness is due to

Lemma 7. *We have $P_{w\gamma}(M) = M \cap wP_\gamma w^{-1}$.*

Proof The inclusion $P_{w\gamma}(M) \subset M \cap wP_\gamma w^{-1}$ follows from definitions. Further, $M \cap wP_\gamma w^{-1}$ contains the opposite Borel of M , hence is a parabolic subgroup of M (in particular, connected). The assertion follows now from: for a root $\check{\alpha}$ of M we have $U_{\check{\alpha}} \subset P_{w\gamma}(M)$ if and only if $U_{\check{\alpha}} \subset M \cap wP_\gamma w^{-1}$. $\square$

We see that $U(P)$ acts transitively on the fibres of (13). So, (13) is a fibration with fibre isomorphic to an affine space of dimension $\langle \gamma + w\gamma, \check{\rho} \rangle - \langle w\gamma, 2\check{\rho}_M \rangle$.

Remark 2. For $w\gamma \in \Lambda_M^{+,\theta}$ one can calculate the dimension of $\text{Gr}_G^\gamma \cap S_P^{w\gamma}$ as follows. Stratify it by B -orbits, that is, by the schemes $\text{Gr}_G^\gamma \cap S^{w_1 w\gamma}$ with $w_1 \in W_M$. Then $\text{Gr}_G^\gamma \cap S^{w_1 w\gamma}$ is an affine space of dimension $\langle \gamma + w_1 w\gamma, \check{\rho} \rangle$. The maximum of these numbers, as w_1 ranges through W_M is $\langle \gamma + w\gamma, \check{\rho} \rangle$.

4.5 Consider a collection $\tilde{\mu} = (\mu_1, \dots, \mu_d)$ with $\mu = \mu_1 + \dots + \mu_d$ and $\mu_i \in \pi_1^\theta(M)$. Let $\mathcal{H}_M^{+,\tilde{\mu}}$ be the stack of collections

$$(\mathcal{F}_M^1, \dots, \mathcal{F}_M^{d+1}, x_1, \dots, x_d \in X, \beta^i), \quad (14)$$

where $\beta^i : \mathcal{F}_M^i |_{X-x_i} \xrightarrow{\sim} \mathcal{F}_M^{i+1} |_{X-x_i}$ is an isomorphism such that $(\mathcal{F}_M^i, \mathcal{F}_M^{i+1}, \beta^i, x_i) \in \mathcal{H}_M^{+,\mu_i}$ for $i = 1, \dots, d$.

If we denote by λ_i the element of $\Lambda_{M,S}^{+,\theta}$ that maps to μ_i then $\mathcal{F}_M^i$ is in the position λ_i with respect to $\mathcal{F}_M^{i+1}$ at x_i . We have a convolution map

$$\text{conv}^{\tilde{\mu}} : \mathcal{H}_M^{+,\tilde{\mu}} \rightarrow \mathcal{H}_M^{+,\mu}$$

sending (14) to $(\mathcal{F}_M^1, \mathcal{F}_M^{d+1}, \beta, D^\mu)$, where $D^\mu = \sum_i \mu_i x_i$ and $\beta : \mathcal{F}_M^1 |_{X-D} \xrightarrow{\sim} \mathcal{F}_M^{d+1} |_{X-D}$ with $D = s_M(D^\mu)$.

Denote by $s^{\tilde{\mu}} : \mathcal{H}_M^{+,\tilde{\mu}} \rightarrow X^d$ the map sending (14) to $(x_1, \dots, x_d)$. From (Lemma 9.3, [14]) one derives

Lemma 8. *i) The map $\text{conv}^{\tilde{\mu}}$ is representable, proper and small over its image. Besides, the perverse sheaf*

$$\text{conv}_!^{\tilde{\mu}}(s^{\tilde{\mu}})^* W^{\boxtimes d}[a]\left(\frac{a}{2}\right) \quad (15)$$

is the Goresky-MacPherson extension from ${}^{rss}\mathcal{H}_M^{+, \mu}$. Here $a = \dim \mathcal{H}_M^{+, \tilde{\mu}}$.

ii) The d -tuple $\tilde{\mu}$ gives rise to $\mathfrak{A}(\mu)$ in general position, say $\mu = \sum_k n_k \nu_k$. The group $\prod_k S_{n_k}$ acts naturally on (15), and the sheaf of $\prod_k S_{n_k}$ -invariants is canonically isomorphic to the direct summand of $\mathcal{L}_W^\mu$ corresponding to $\mathfrak{A}(\mu)$. $\square$

Remark 3. From Lemma 8 it follows that for any $\mu \in \pi_1^+(M)$ the complex $\mathcal{L}_W^\mu$ is ULA with respect to both projections $\mathfrak{p}_M, \mathfrak{q}_M : \mathcal{H}_M^{+, \mu} \rightarrow \text{Bun}_M$.

Proof of Proposition 8

1) Consider the diagram

$$\begin{array}{ccc} \mathcal{H}_P^{+, \mu} & \xrightarrow{q_M} & \mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P \\ & \searrow \mathfrak{q}_P \times \text{supp}_P & \downarrow \\ & & \text{Bun}_P \times X_M^\mu \end{array}$$

For any k -point of $\text{Bun}_P \times X_M^\mu$ given by $(\mathcal{F}'_P, D^\mu = \sum_k \mu_k x_k)$, where x_k are pairwise distinct, over $(\mathcal{F}'_P, D^\mu = \sum_k \mu_k x_k)$ the map q_M becomes the product

$$\prod_k \text{Gr}_G^+ \cap S_P^{\mu_k} \rightarrow \prod_k \text{Gr}_M^{+, \mu_k}$$

of the maps t_S^{+, μ_k} . In particular, q_M is surjective.

Let $\mathfrak{A}(\mu)$ in general position be given by $\mu = \sum_k n_k \mu_k$. Let $U \subset {}^{rss}\mathcal{H}_M^{+, \mu}$ be the preimage of the corresponding connected component of ${}^{rss}X_M^\mu$. Over the open substack $U \times_{\text{Bun}_M} \text{Bun}_P$, the map q_M is a fibration with fibre isomorphic to an affine space of dimension

$$a(\mathfrak{A}(\mu)) \stackrel{\text{def}}{=} \langle d\gamma, \check{\rho} \rangle + \sum_k \langle n_k \lambda_k, \check{\rho} - 2\check{\rho}_M \rangle,$$

where $\lambda_k \in \Lambda_{M,S}^{+, \theta}$ maps to $\mu_k \in \pi_1^\theta(M)$.

The restriction of $f_M^* \mathcal{L}_W^d$ to $q_M^*(U \times_{\text{Bun}_M} \text{Bun}_P)$ comes from ${}^{rss}X^{(d)}$. So, over $U \times_{\text{Bun}_M} \text{Bun}_P$, we get the desired isomorphism. Now it suffices to show that, up to a shift, $q_M! f_M^* \mathcal{S}pr_W^d$ is a perverse sheaf, the Goresky-MacPherson extension from ${}^{rss}\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P$.

For a d -tuple $\tilde{\mu} = (\mu_1, \dots, \mu_d)$ with $\mu = \mu_1 + \dots + \mu_d$ and $\mu_i \in \pi_1^\theta(M)$, let $\mathcal{H}_P^{+, \tilde{\mu}}$ be the stack of collections

$$(\mathcal{F}_P^1, \dots, \mathcal{F}_P^{d+1}, x_1, \dots, x_d, \beta^i), \quad (16)$$

where $x_i \in X$ and $(\mathcal{F}_P^i, \mathcal{F}_P^{i+1}, x_i, \beta^i) \in \mathcal{H}_P^{+, \mu_i}$ for $i = 1, \dots, d$. The stack

$$\mathcal{H}_P^{+, \mu} \times_{(\mathcal{H}_G^{+, d} \times_{\text{Bun}_G} \text{Bun}_P)} (\tilde{\mathcal{H}}_G^{+, d} \times_{\text{Bun}_G} \text{Bun}_P)$$

is stratified by locally closed substacks $\mathcal{H}_P^{+, \tilde{\mu}}$ indexed by such tuples $\tilde{\mu}$.

We have a diagram

$$X^{(d)} \xleftarrow{s^{\tilde{\mu}}} \mathcal{H}_P^{+, \tilde{\mu}} \xrightarrow{\alpha} \mathcal{H}_M^{+, \tilde{\mu}} \times_{\text{Bun}_M} \text{Bun}_P \rightarrow \mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P,$$

where α sends (16) to $(\mathcal{F}_M^1, \dots, \mathcal{F}_M^d, \mathcal{F}_P^{d+1}, x_1, \dots, x_d, \beta^i)$ with $\mathcal{F}_M^i = \mathcal{F}_P^i \times_P M$, and the last map is $\text{conv}^{\tilde{\mu}} \times \text{id}$. It suffices to show that for each $\tilde{\mu}$ as above,

$$(\text{conv}^{\tilde{\mu}} \times \text{id})_! \alpha_! (s^{\tilde{\mu}})^* W^{(d)}$$

is a perverse sheaf (up to a shift), the Goresky-MacPherson extension from ${}^{rss}\mathcal{H}_M^{+, \mu} \times_{\text{Bun}_M} \text{Bun}_P$. Since α is a composition of affine fibrations, our statement about $q_M! f_M^* \mathcal{L}_W^d$ follows from Lemma 8, i).

2) Applying 1) to the 1-admissible data $\{-w_0(\gamma)\}$, one gets the formula for $p_M! f_M^* \mathcal{L}_W^d$. $\square$

4.6.1 Averaging functors for Levi subgroups

Recall for any $\mu \in \pi_1^+(M)$ the diagram $\text{Bun}_M \xleftarrow{\mathfrak{p}_M} \mathcal{H}_M^{+, \mu} \xrightarrow{\mathfrak{q}_M} \text{Bun}_M$. For a local system W on X denote by $\text{Av}_W^\mu : \text{D}(\text{Bun}_M) \rightarrow \text{D}(\text{Bun}_M)$ the functor

$$\text{Av}_W^\mu(K) = (\mathfrak{q}_M)_! (\mathfrak{p}_M^* K \otimes \mathcal{L}_W^\mu)[-d_M] \left(\frac{-d_M}{2}\right)$$

Let also $\text{Av}_W^{-\mu} : \text{D}(\text{Bun}_M) \rightarrow \text{D}(\text{Bun}_M)$ be given by

$$\text{Av}_W^{-\mu}(K) = (\mathfrak{p}_M)_! (\mathfrak{q}_M^* K \otimes \mathcal{L}_W^\mu)[-d_M] \left(\frac{-d_M}{2}\right)$$

By Remark 3, we have $\mathbb{D} \circ \text{Av}_W^\mu \xrightarrow{\sim} \text{Av}_{W^*}^\mu \circ \mathbb{D}$ and $\mathbb{D} \circ \text{Av}_W^{-\mu} \xrightarrow{\sim} \text{Av}_{W^*}^{-\mu} \circ \mathbb{D}$ naturally.

The proof of the following result is completely analogous to that of Proposition 3.

Proposition 9. *Let $\mu \in \pi_1^+(M)$. Let $\mathfrak{A}(\mu)$ in general position be given by $\mu = \sum_k n_k \mu_k$. Recall the map $s^{\mathfrak{A}(\mu)} : X^{\mathfrak{A}(\mu)} \rightarrow X_M^\mu$ (cf. sect. 4.3). Let $\lambda_k \in \Lambda_{M,S}^{+, \theta}$ be the element that maps to $\mu_k \in \pi_1^\theta(M)$. Let W be any local system on X . Let E be a $\check{M}$ -local system on X , K be E -Hecke eigensheaf on Bun_M . Then for the diagram*

$$\text{Bun}_M \xleftarrow{\mathfrak{p}_M} \mathcal{H}_M^{+, \mu} \xrightarrow{\mathfrak{q}_M \times \text{supp}_M} \text{Bun}_M \times X_M^\mu$$

there is a canonical isomorphism

$$(\mathfrak{q}_M \times \text{supp}_M)_! (\mathfrak{p}_M^* K \otimes \mathcal{L}_W^{\mathfrak{A}(\mu)})[-d_M] \left(\frac{-d_M}{2}\right) \xrightarrow{\sim} K \boxtimes s_!^{\mathfrak{A}(\mu)} \left(\boxtimes_k (W \otimes U_E^{\lambda_k})^{(n_k)} \right) [d] \left(\frac{d}{2}\right) \quad \square$$

Corollary 3. *For each standard proper parabolic P of G there exists a constant $c(P)$ with the following property. Let E be any $\check{M}$ -local system E on X , K be a E -Hecke eigensheaf on Bun_M . Let W be an irreducible local system on X of rank $r = \dim V^\gamma$. For any $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is $d\theta$ with $d > c(P)$ we have $\text{Av}_W^\mu(K) = 0$.*

Proof The functor $\mathrm{Av}_W^\mu = \oplus \mathrm{Av}_W^{\mathfrak{A}(\mu)}$ is a direct sum of functors indexed by $\mathfrak{A}(\mu)$ in general position. In the notation of Proposition 9, we have

$$\mathrm{Av}_W^{\mathfrak{A}(\mu)}(K) \xrightarrow{\sim} K \otimes (\otimes_k \mathrm{Sym}^{n_k} \mathrm{R}\Gamma(X, W \otimes U_E^{\lambda_k})) [d] \left(\frac{d}{2}\right) \quad (17)$$

Here $d = \sum n_k$, and k runs through the finite set $\pi_1^\theta(M)$. For d large enough at least one of n_k will satisfy $n_k > r(2g - 2) \dim U^{\lambda_k}$, and the RHS of (17) will vanish. $\square$

Generalizing the Vanishing Conjecture of Frenkel, Gaitsgory and Vilonen ([4]), we suggest

Conjecture 4. *Let W be an irreducible local system on X of rank $r = \dim V^\gamma$. Assume that P is a standard proper parabolic of G . Then for all $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ equals $d\theta$ with $d > c(P)$, the functor Av_W^μ vanishes identically.*

4.6.2 Consider the diagram $\mathrm{Bun}_G \xleftarrow{\alpha_P} \mathrm{Bun}_P \xrightarrow{\beta_P} \mathrm{Bun}_M$, where α_P and β_P are natural maps. The constant term functor $\mathrm{CT}_P : \mathrm{D}(\mathrm{Bun}_G) \rightarrow \mathrm{D}(\mathrm{Bun}_M)$ is defined by $\mathrm{CT}_P(K) = \beta_{P!} \alpha_P^*(K)$.

The following is a generalization of Lemma 9.8, [4].

Lemma 9. *Let W be any local system on X . For any $K \in \mathrm{D}(\mathrm{Bun}_G)$ and $d \geq 0$ the complex $\mathrm{CT}_P \circ \mathrm{Av}_W^d(K) \in \mathrm{D}(\mathrm{Bun}_M)$ has a canonical filtration by complexes*

$$\mathrm{Av}_W^\mu \circ \mathrm{CT}_P(K) \otimes \bar{\mathbb{Q}}_\ell[1] \left(\frac{1}{2}\right)^{\otimes \langle \mu, 2\check{\rho} - 2\check{\rho}_M \rangle} \quad (18)$$

indexed by those $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is $d\theta$.

Proof Consider the stack $\mathcal{H}_G^{+,d} \times_{\mathrm{Bun}_G} \mathrm{Bun}_P$, where we used $\mathfrak{q} : \mathcal{H}_G^{+,d} \rightarrow \mathrm{Bun}_G$ in the fibred product. The complex $\mathrm{CT}_P \circ \mathrm{Av}_W^d(K)$ is the direct image with respect to the natural map

$$\mathcal{H}_G^{+,d} \times_{\mathrm{Bun}_G} \mathrm{Bun}_P \rightarrow \mathrm{Bun}_M \quad (19)$$

Recall that $\mathcal{H}_G^{+,d} \times_{\mathrm{Bun}_G} \mathrm{Bun}_P$ is stratified by locally closed substacks $\mathcal{H}_P^{+,\mu}$ indexed by those $\mu \in \pi_1^+(M)$ whose image in $\pi_1(G)$ is $d\theta$. This gives a filtration on $\mathrm{CT}_P \circ \mathrm{Av}_W^d(K)$.

The restriction of the map (19) to the stratum $\mathcal{H}_P^{+,\mu}$ can be written as a composition

$$\mathcal{H}_P^{+,\mu} \xrightarrow{p_M} \mathcal{H}_M^{+,\mu} \times_{\mathrm{Bun}_M} \mathrm{Bun}_P \rightarrow \mathrm{Bun}_M$$

So, by Proposition 8, the contribution of the stratum $\mathcal{H}_P^{+,\mu}$ to the direct image in question is exactly (18). $\square$

Corollary 4. *Assume that Conjecture 4 holds. Then*

- 1) *Let d satisfy $d > c(P)$ for any standard proper parabolic of G . Then for any $K \in \mathrm{D}(\mathrm{Bun}_G)$ and any irreducible local system W on X of rank $r = \dim V^\gamma$ the complex $\mathrm{Av}_W^d(K)$ is cuspidal.*
- 2) *Let E be $\check{G}$ -local system on X and K be a E -Hecke eigensheaf on Bun_G . If V_E^γ is irreducible then K is cuspidal.*

Proof 1) is clear.

2) The argument given in ([4], Theorem 9.2) applies in our setting. Namely, pick d such that $d > c(P)$ for any standard proper parabolic of G . Set $W = (V_E^\gamma)^*$. By Proposition 3,

$$\mathrm{CT}_P \circ \mathrm{Av}_W^d(K) \xrightarrow{\sim} \mathrm{CT}_P(K) \otimes \mathrm{R}\Gamma(X^{(d)}, (W \otimes V_E^\gamma)^{(d)})[d](\frac{d}{2})$$

The LHS vanishes by Lemma 9. Since $\mathrm{R}\Gamma(X^{(d)}, (W \otimes V_E^\gamma)^{(d)})$ is not zero, $\mathrm{CT}_P(K) = 0$. $\square$

Remark 4. For $G = \mathrm{GL}_n$ Conjecture 4 is proved by D. Gaitsgory ([5]). For $G = \mathrm{GSp}_4$ (example 2 in the appendix) Conjecture 4 also holds, it is easily reduced to the result of *loc.cit.* for GL_2 . So, for $G = \mathrm{GSp}_4$ Corolary 4 is unconditional.

APPENDIX. 1-ADMISSIBLE GROUPS

Definition 8. Let H be a connected, semi-simple and simply-connected group (over k). Assume that the center $Z(H)$ is cyclic of order h and fix an isomorphism $\mathfrak{z} : \mu_h \xrightarrow{\sim} Z(H)$. Assume that the characteristic of k does not divide h . Denote by G the quotient of $H \times \mathbb{G}_m$ by the diagonally embedded μ_h . Call a reductive group G over k *1-admissible*, if it is obtained in this way.

Let H be a connected, semi-simple and simply-connected group (over k). Let T_H be a maximal torus of H . Write $\check{\Lambda}_H$ (resp., Λ_H) for the weight (resp., coweight) lattice of T_H . Let $\check{Q}_H \subset \check{\Lambda}_H$ be the root lattice. Set

$$Q_H = \{\lambda \in \Lambda_H \otimes \mathbb{Q} \mid \langle \lambda, \check{\lambda} \rangle \in \mathbb{Z} \text{ for any } \check{\lambda} \in \check{Q}_H\}$$

We have a natural pairing $\check{\Lambda}_H / \check{Q}_H \times Q_H / \Lambda_H \rightarrow (\frac{1}{h}\mathbb{Z}) / \mathbb{Z}$. Therefore, any isomorphism

$$\tau : Q_H / \Lambda_H \xrightarrow{\sim} (\frac{1}{h}\mathbb{Z}) / \mathbb{Z}$$

yields an isomorphism $\check{\tau} : \check{\Lambda}_H / \check{Q}_H \xrightarrow{\sim} \mathbb{Z} / h\mathbb{Z}$. Since the characteristic of k does not divide h , $\mathrm{Hom}(Z(H), k^*) \xrightarrow{\sim} \check{\Lambda}_H / \check{Q}_H$ canonically. So, $\check{\tau}$ yields $\mathfrak{z} : \mu_h \xrightarrow{\sim} Z(H)$ such that for $x \in \mu_h$, $\check{\lambda} \in \check{\Lambda}_H$ we have

$$\check{\lambda}(\mathfrak{z}(x)) = x^{\check{\tau}(\check{\lambda})}$$

So, τ gives rise to a 1-admissible group $G = (H \times \mathbb{G}_m) / \mu_h$.

For the maximal torus $T = (T_H \times \mathbb{G}_m) / \mu_h$ in G the weight lattice is

$$\check{\Lambda} = \{(\check{\lambda}, a) \in \check{\Lambda}_H \times \mathbb{Z} \mid \check{\tau}(\check{\lambda}) + a = 0 \pmod{h}\}$$

and the coweight lattice is

$$\Lambda = \{(\lambda, b) \in Q_H \times (\frac{1}{h}\mathbb{Z}) \mid \tau(\lambda) - b \in \mathbb{Z}\}$$

It is understood that the pairing $\Lambda \times \check{\Lambda} \rightarrow \mathbb{Z}$ sends $(\lambda, b), (\check{\lambda}, a)$ to $\langle \lambda, \check{\lambda} \rangle + ab$. The map $(\lambda, b) \mapsto b$ yields an isomorphism $\pi_1(G) \xrightarrow{\sim} \Lambda / \Lambda_H \xrightarrow{\sim} \frac{1}{h}\mathbb{Z}$. Note also that $\pi_1(\check{G}) \xrightarrow{\sim} \check{\Lambda} / \check{Q}_H \xrightarrow{\sim} \mathbb{Z}$.

The next result follows from definitions.

Lemma 10. Let $\gamma_H \in Q_H$ be a dominant coweight for $H_{ad} = H/Z(H)$. Assume that

- either γ_H is minuscule or ($H = 1$ and $\gamma_H = 0$);
- γ_H generates Q_H/Λ_H ;
- the irreducible representation V^{γ_H} of $(H_{ad})^\vee$ is faithful.

Let $\tau : Q_H/\Lambda_H \xrightarrow{\sim} (\frac{1}{h}\mathbb{Z})/\mathbb{Z}$ be the isomorphism sending γ_H to $\frac{1}{h}$, and G be the corresponding 1-admissible group. Set $\gamma = (\gamma_H, \frac{1}{h}) \in \Lambda$. Then $\{\gamma\}$ is a 1-admissible datum for G . $\square$

EXAMPLES OF 1-ADMISSIBLE DATA

The examples below are produced using Lemma 10.

1. **The case $G = \mathrm{GL}_n$.** In the standard notation $\Lambda = \mathbb{Z}^n$, $\check{\Lambda} = \mathbb{Z}^n$. For $1 \leq i < n$ take $\check{\omega}_i = (1, \dots, 1, 0, \dots, 0)$ where 1 appears i times, and $\check{\omega}_0 = (1, \dots, 1)$. All the conditions are satisfied and $\gamma = (1, 0, \dots, 0)$.

Let $\gamma_i = (1, \dots, 1, 0, \dots, 0) \in \Lambda$, where 1 appears i times. Then $\Lambda_{G,S}^+$ is the $\mathbb{Z}_+$ -span of $\gamma_1, \gamma_2, \dots, \gamma_n$. So, $\Lambda_{G,S}^+ \xrightarrow{\sim} (\mathbb{Z}_+)^n$. The element $\omega = \gamma_n$ generates the group of coweights orthogonal to all roots.

Remark 5. This particular choice of γ_H yields a construction of an automorphic sheaf proposed by Laumon in [9]. However, all fundamental coweights for $H_{ad} = \mathrm{PSL}_n$ are minuscule, and a choice of γ_H here is equivalent to a choice of a generator of the cyclic group $\pi_1(H_{ad})$. If γ_H is a fundamental coweight corresponding to a simple root which is not one of two edges of the Dynkin diagram A_{n-1} then the corresponding 1-admissible group G is not isomorphic to GL_n .

2. **The case $G = G\mathrm{Sp}_{2n}$, $n \geq 1$.** The group G is a quotient of $\mathbb{G}_m \times \mathrm{Sp}_{2n}$ by the diagonally embedded $\{\pm 1\}$. Realise G as the subgroup of $\mathrm{GL}(k^{2n})$ preserving up to a scalar the bilinear form given by the matrix

$$\begin{pmatrix} 0 & E_n \\ -E_n & 0 \end{pmatrix},$$

where E_n is the unit matrix of GL_n .

The maximal torus T of G is $\{(y_1, \dots, y_{2n}) \mid y_i y_{n+i} \text{ does not depend on } i\}$. Let $\check{\epsilon}_i \in \check{\Lambda}$ be the character that sends a point of T to y_i . The roots are

$$\check{R} = \{\pm \check{\alpha}_{ij} \ (i < j \in 1, \dots, n), \pm \check{\beta}_{ij} \ (i \leq j \in 1, \dots, n)\},$$

where $\check{\alpha}_{ij} = \check{\epsilon}_i - \check{\epsilon}_j$ and $\check{\beta}_{ij} = \check{\epsilon}_i - \check{\epsilon}_{n+j}$.

We have $\Lambda = \{(a_1, \dots, a_{2n}) \mid a_i + a_{n+i} \text{ does not depend on } i\}$. The weight lattice is

$$\check{\Lambda} = \mathbb{Z}^{2n} / \{\check{\epsilon}_i + \check{\epsilon}_{n+i} - \check{\epsilon}_j - \check{\epsilon}_{n+j}, \ i < j\}$$

Let e_i denote the standard basis of $\mathbb{Z}^{2n}$. The coroots are

$$R = \{\pm\alpha_{ij} \ (i < j \in 1, \dots, n), \pm\beta_{ij} \ (i \leq j \in 1, \dots, n)\},$$

where $\beta_{ij} = e_i + e_j - e_{n+i} - e_{n+j}$ for $i < j$ and $\beta_{ii} = e_i - e_{n+i}$. Besides, $\alpha_{ij} = e_i + e_{n+j} - e_j - e_{n+i}$.

Fix positive roots

$$\check{R}^+ = \{\check{\alpha}_{ij} \ (i < j \in 1, \dots, n), \check{\beta}_{ij} \ (i \leq j \in 1, \dots, n)\}$$

Then the simple roots are $\check{\alpha}_{12}, \dots, \check{\alpha}_{n-1,n}$ and $\check{\beta}_{n,n}$.

For $1 \leq i < n$ pick the fundamental weight $\check{\omega}_i$ corresponding to the simple coroot $\alpha_{i,i+1}$ to be $\check{\omega}_i = (1, \dots, 1, 0, \dots, 0)$, where 1 appears i times, and 0 appears $2n - i$ times. Let the fundamental weight $\check{\omega}_n$ corresponding to $\beta_{n,n}$ be $\check{\omega}_n = (1, \dots, 1, 0, \dots, 0)$, where 1 appears n times. The orthogonal to the coroot lattice is the subgroup $\mathbb{Z}\check{\omega}_0$ with $\check{\omega}_0 = (1, 0, \dots, 0; 1, 0, \dots, 0)$.

All our conditions are satisfied and $\gamma = (1, \dots, 1; 0, \dots, 0)$ (here 1 appears n times).

For $1 \leq i < n$ let $\gamma_i = (2, \dots, 2, 1, \dots, 1; 0, \dots, 0, 1, \dots, 1)$, where 2 appears i times then 1 appears $n - i$ times then 0 appears i times and finally 1 appears $n - i$ times. The element $\omega = (1, \dots, 1) \in \Lambda$ generates the group of coweights orthogonal to all roots. The semigroup $\Lambda_{G,S}^+$ is the $\mathbb{Z}_+$ -span of $\gamma, \gamma_1, \dots, \gamma_{n-1}, \omega$. In fact, these $n + 1$ elements are linearly independent in Λ , so $\Lambda_{G,S}^+ \xrightarrow{\sim} (\mathbb{Z}_+)^{n+1}$.

Note that V^γ is the spinor representation of $\check{G} \xrightarrow{\sim} \mathrm{GSpin}_{2n+1}$ of dimension 2^n . We have

$$V^\gamma \otimes V^\gamma \xrightarrow{\sim} V^{2\gamma} \oplus V^\omega \oplus \sum_{i=1}^{n-1} V^{\gamma_i}$$

and $\dim V^{\gamma_1} = 2n + 1$. Besides, $\wedge^i V^{\gamma_1} \xrightarrow{\sim} V^{\gamma_1 + (i-1)\omega}$ for $i = 1, \dots, n-1$ and $\wedge^n V^{\gamma_1} \xrightarrow{\sim} V^{2\gamma_1 + (n-1)\omega}$. There is an exact sequence $1 \rightarrow \mathbb{G}_m \rightarrow \check{G} \rightarrow \mathrm{SO}_{2n+1} \rightarrow 1$, and $V^{\gamma_1 - \omega}$ comes from the standard representation of SO_{2n+1} .

3. The case $G = \mathrm{GSpin}_{2n+1}$, $n \geq 1$. The group G is the quotient of $\mathbb{G}_m \times \mathrm{Spin}_{2n+1}$ by the diagonally embedded $\{\pm 1\}$. We have $\check{G} \xrightarrow{\sim} \mathrm{GSp}_{2n}$, the root data for G is dual to that of example 2. Interchanging the role of objects and coobjects in example 2, we get

$$\Lambda = \mathbb{Z}^{2n} / \{\epsilon_i + \epsilon_{n+i} - \epsilon_j - \epsilon_{n+j}, \ i < j\}$$

where (ϵ_i) is the standard basis of $\mathbb{Z}^{2n}$. The weight lattice is

$$\check{\Lambda} = \{(a_1, \dots, a_{2n}) \in \mathbb{Z}^{2n} \mid a_i + a_{n+i} \text{ does not depend on } i\}$$

Define $\check{\omega}_i$ as follows. For $1 \leq i < n$ let

$$\check{\omega}_i = (2, \dots, 2, 1, \dots, 1; 0, \dots, 0, 1, \dots, 1) \in \check{\Lambda}$$

where 2 appears i times then 1 appears $n - i$ times then 0 appears i times and finally 1 appears $n - i$ times. Let $\check{\omega}_n = (1, \dots, 1; 0, \dots, 0)$ where 1 appears n times. Let $\check{\omega}_0 = (1, \dots, 1)$.

All the conditions are satisfied and $\gamma = (1, 0, \dots, 0) \in \Lambda$.

For $1 \leq i \leq n$ let $\gamma_i = (1, \dots, 1, 0, \dots, 0) \in \Lambda$, where 1 appears i times and 0 appears $2n - i$ times. The group of coweights orthogonal to all roots is $\mathbb{Z}\omega$ with $\omega = (1, 0, \dots, 0; 1, 0, \dots, 0)$, here 1 appears on the first and $(n+1)$ -th places. The semigroup $\Lambda_{G,S}^+$ is the $\mathbb{Z}_+$ -span of $\gamma_1, \dots, \gamma_n, \omega$. These $n+1$ elements are linearly independent in Λ , so $\Lambda_{G,S}^+ \xrightarrow{\sim} (\mathbb{Z}_+)^{n+1}$.

We have $\wedge^2 V^\gamma = V^{\gamma_2} \oplus V^\omega$. To assure condition (A) of Conjecture 1, it suffices to require that V_E^γ is irreducible and $V_E^{\gamma_2}$ has no local subsystems of rank one.

4. The case $H = \text{Spin}_{2n}$, $n \geq 3$ odd. We have

$$\check{\Lambda}_H = \{(a_1, \dots, a_n) \in (\frac{1}{2}\mathbb{Z})^n \mid a_i - a_j \in \mathbb{Z}\}$$

and $\Lambda_H = \{(a_1, \dots, a_n) \in \mathbb{Z}^n \mid a_1 + \dots + a_n \text{ is even}\}$. The roots are

$$\check{R} = \{\pm\check{\alpha}_{ij}, \pm\check{\beta}_{ij} \mid (1 \leq i < j \leq n)\}$$

with $\check{\alpha}_{ij} = \check{e}_i - \check{e}_j$ and $\check{\beta}_{ij} = \check{e}_i + \check{e}_j$. The coroots are

$$R = \{\pm\alpha_{ij}, \pm\beta_{ij} \mid (1 \leq i < j \leq n)\}$$

with $\alpha_{ij} = e_i - e_j$ and $\beta_{ij} = e_i + e_j$. Pick positive roots $\check{R} = \{\check{\alpha}_{ij}, \check{\beta}_{ij} \mid (1 \leq i < j \leq n)\}$. Then simple roots are

$$\check{\alpha}_{12}, \dots, \check{\alpha}_{n-1,n}, \check{\beta}_{n-1,n}$$

We have $\check{Q}_H = \{(a_1, \dots, a_n) \in \mathbb{Z}^n \mid a_1 + \dots + a_n \text{ is even}\}$,

$$Q_H = \{(a_1, \dots, a_n) \in (\frac{1}{2}\mathbb{Z})^n \mid a_i - a_j \in \mathbb{Z}\}$$

There are two possible choices for γ_H , namely $\gamma_H = (\frac{1}{2}, \dots, \frac{1}{2})$ or $\gamma_H = (\frac{1}{2}, \dots, \frac{1}{2}, -\frac{1}{2})$, the fundamental coweights corresponding to $\check{\beta}_{n-1,n}$ or to $\check{\alpha}_{n-1,n}$ respectively.

5. The case E_6 . So, H is the simply-connected group corresponding to E_6 root system. There are two possible choices for γ_H , namely ω_1 or ω_6 , the fundamental coweights corresponding to the simple roots $\check{\alpha}_1$ and $\check{\alpha}_6$ in the Bourbaki table ([1], Ch. 6, no. 4.12, p.219).

6. The case E_7 . So, H is the simply-connected group corresponding to E_7 root system. Take γ_H to be the fundamental coweight ω_7 corresponding to the root $\check{\alpha}_7$ in the Bourbaki table ([1], Ch. 6, no. 4.11, p. 217).

References

- [1] N. Bourbaki, Groupes et algèbres de Lie, Éléments de Mathématique, chapitres 4,5 et 6, Masson (1981)

- [2] A. Braverman, D. Gaitsgory, Geometric Eisenstein series, math.AG/9912097
- [3] E. Frenkel, D. Gaitsgory, K. Vilonen, Whittaker patterns in the geometry of moduli spaces of bundles on curves. *Ann. of Math.* (2) 153 (2001), no. 3, 699–748.
- [4] E. Frenkel, D. Gaitsgory, K. Vilonen, On the geometric Langlands conjecture. *J. Amer. Math. Soc.* 15 (2002), no. 2, 367–417
- [5] D. Gaitsgory, On a vanishing conjecture appearing in the geometric Langlands correspondence, math.AG/0204081
- [6] D. Ginzburg, S. Rallis, D. Soudry, Self-dual automorphic GL_n -modules and construction of a backward lifting from GL_n to classical groups, *IMRN*, no.14 (1997), p. 687-701.
- [7] D. Ginzburg, S. Rallis, D. Soudry, On explicit lifts of cusp forms from GL_m to classical groups, *Ann. of Math.* 150 (1999), p. 807-866.
- [8] L. Lafforgue, Chtoucas de Drinfeld et correspondance de Langlands, prépublication 2000-62, Université Paris-Sud.
- [9] G. Laumon, Correspondance de Langlands géométrique pour les corps de fonctions, *Duke Math. J.*, vol. 54, No. 2 (1987) p. 309-359
- [10] G. Laumon, Faisceaux automorphes pour $GL(n)$: la première construction de Drinfeld, alg-geom/9511004
- [11] S. Lysenko, Local geometrized Rankin-Selberg method for $GL(n)$, *Duke Math. J.* 111 (2002), no. 3, 451–493.
- [12] S. Lysenko, Orthogonality relations between automorphic sheaves attached to irreducible local systems of rank 2 on a curve, PhD thesis, Université Paris-Sud (1999)
- [13] C. Mœglin, J.-L. Waldspurger, Spectral decomposition and Eisenstein series. Une paraphrase de l'Écriture. *Cambridge Tracts in Mathematics*, 113. Cambridge University Press, Cambridge, 1995.
- [14] B.C. Ngo, P. Polo, Résolutions de Demazure affines et formule de Casselman-Shalika géométrique, mathAG/0005022
- [15] D. Soudry, Rankin-Selberg convolutions for $SO_{2l+1} \times GL_n$: local theory, *Memoirs AMS* no. 500, v. 105 (1993)